ARCHIVUM MATHEMATICUM (BRNO)
Tomus 47 (2011), 201–215

INDUCED DIFFERENTIAL FORMS ON MANIFOLDS OF FUNCTIONS

Cornelia Vizman

Dedicated to Peter W. Michor at the occasion of his 60th birthday

Abstract. Differential forms on the Fréchet manifold $\mathcal{F}(S,M)$ of smooth functions on a compact k-dimensional manifold S can be obtained in a natural way from pairs of differential forms on M and S by the hat pairing. Special cases are the transgression map $\Omega^p(M) \to \Omega^{p-k}(\mathcal{F}(S,M))$ (hat pairing with a constant function) and the bar map $\Omega^p(M) \to \Omega^p(\mathcal{F}(S,M))$ (hat pairing with a volume form). We develop a hat calculus similar to the tilda calculus for non-linear Grassmannians [6].

1. Introduction

Pairs of differential forms on the finite dimensional manifolds M and S induce differential forms on the Fréchet manifold $\mathcal{F}(S,M)$ of smooth functions. More precisely, if S is a compact oriented k-dimensional manifold, the hat pairing is:

$$\Omega^p(M) \times \Omega^q(S) \to \Omega^{p+q-k}(\mathcal{F}(S,M))$$
$$\widehat{\omega \cdot \alpha} = \int\!\!\!\!\!\!- _S \operatorname{ev}^* \omega \wedge \operatorname{pr}^* \alpha \,,$$

where $\operatorname{ev}\colon S \times \mathcal{F}(S,M) \to M$ denotes the evaluation map, $\operatorname{pr}\colon S \times \mathcal{F}(S,M) \to S$ the projection and $\int\!\!\!\!\!\!-_S$ fiber integration. We show that the hat pairing is compatible with the canonical $\mathrm{Diff(M)}$ and $\mathrm{Diff}(S)$ actions on $\mathcal{F}(S,M)$, and with the exterior derivative. As a consequence we obtain a hat pairing in cohomology.

The hat (transgression) map is the hat pairing with the constant function 1, so it associates to any form $\omega \in \Omega^p(M)$ the form $\widehat{\omega \cdot 1} = \widehat{\omega} = \int\!\!\!\!\!\!-_S \operatorname{ev}^* \omega \in \Omega^{p-k}(\mathcal{F}(S,M))$. Since $\mathfrak{X}(M)$ acts infinitesimally transitive on the open subset $\mathrm{Emb}(S,M) \subset \mathcal{F}(S,M)$ of embeddings of the k-dimensional oriented manifold S into M [7], the expression of $\widehat{\omega}$ at $f \in \mathrm{Emb}(S,M)$ is

$$\widehat{\omega}(X_1 \circ f, \dots, X_{p-k} \circ f) = \int_S f^*(i_{X_{p-k}} \dots i_{X_1} \omega), \quad X_1, \dots, X_{p-k} \in \mathfrak{X}(M)\,.$$

2010 *Mathematics Subject Classification*: primary 11K11; secondary 22C22.
Key words and phrases: manifold of functions, fiber integral, diffeomorphism group.
Received January 18, 2011, revised March 2011. Editor J. Slovák.

When S is the circle, then one obtains the usual transgression map with values in the space of $(p-1)$-forms on the free loop space of M.

Let $\mathrm{Gr}_k(M)$ be the non-linear Grassmannian of k-dimensional oriented submanifolds of M. The tilda map associates to every $\omega \in \Omega^p(M)$ a differential $(p-k)$-form on $\mathrm{Gr}_k(M)$ given by [6]

$$\tilde{\omega}(\tilde{Y}_N^1, \dots, \tilde{Y}_N^{p-k}) = \int_N i_{Y_N^{p-k}} \cdots i_{Y_N^1} \omega\,, \quad \forall \tilde{Y}_N^1, \dots, \tilde{Y}_N^{p-k} \in \Gamma(TN^\perp) = T_N \,\mathrm{Gr}_k(M)\,,$$

for $\tilde{Y}_N$ section of the orthogonal bundle $TN^\perp$ represented by the section Y_N of $TM|_N$. The natural map

$$\pi\colon \mathrm{Emb}(S, M) \to \mathrm{Gr}_k(M)\,, \quad \pi(f) = f(S)$$

provides a principal bundle with the group $\mathrm{Diff}_+(S)$ of orientation preserving diffeomorphisms of S as structure group.

The hat map on $\mathrm{Emb}(S, M)$ and the tilda map on $\mathrm{Gr}_k(M)$ are related by $\widehat{\omega} = \pi^* \tilde{\omega}$. This is the reason why for the hat calculus one has similar properties to those for the tilda calculus. The tilda calculus was used to study the non-linear Grassmannian of co-dimension two submanifolds as symplectic manifold [6]. We apply the hat calculus to the hamiltonian formalism for p-branes and open p-branes [1] [2].

The bar map $\bar{\omega} = \widehat{\omega \cdot \mu}$ is the hat pairing with a fixed volume form μ on S, so

$$\bar{\omega}(Y_f^1, \dots, Y_f^p) = \int_S \omega(Y_f^1, \dots, Y_f^p)\mu\,, \quad \forall Y_f^1, \dots, Y_f^p \in \Gamma(f^*TM) = T_f \mathcal{F}(S, M)\,.$$

We use the bar calculus to study $\mathcal{F}(S, M)$ with symplectic form $\bar{\omega}$ induced by a symplectic form ω on M. The natural actions of $\mathrm{Diff}_{\mathrm{ham}}(M, \omega)$ and $\mathrm{Diff}_{\mathrm{ex}}(S, \mu)$, the group of hamiltonian diffeomorphisms of M and the group of exact volume preserving diffeomorphisms of S, are two commuting hamiltonian actions on $\mathcal{F}(S, M)$. Their momentum maps form the dual pair for ideal incompressible fluid flow [12] [4].

We are grateful to Stefan Haller for extremely helpful suggestions.

2. Hat pairing

We denote by $\mathcal{F}(S, M)$ the set of smooth functions from a compact oriented k-dimensional manifold S to a manifold M. It is a Fréchet manifold in a natural way [10]. Tangent vectors at $f \in \mathcal{F}(S, M)$ are identified with vector fields on M along f, i.e. sections of the pull-back vector bundle f^*TM.

Let $\mathrm{ev}\colon S \times \mathcal{F}(S, M) \to M$ be the evaluation map $\mathrm{ev}(x, f) = f(x)$ and $\mathrm{pr}\colon S \times \mathcal{F}(S, M) \to S$ the projection $\mathrm{pr}(x, f) = x$. A pair of differential forms $\omega \in \Omega^p(M)$ and $\alpha \in \Omega^q(S)$ determines a differential form $\widehat{\omega \cdot \alpha}$ on $\mathcal{F}(S, M)$ by the fiber integral over S (whose definition and properties are listed in the appendix) of the $(p+q)$-form $\mathrm{ev}^*\,\omega \wedge \mathrm{pr}^*\,\alpha$ on $S \times \mathcal{F}(S, M)$:

$$\widehat{\omega \cdot \alpha} = \fint_S \mathrm{ev}^*\,\omega \wedge \mathrm{pr}^*\,\alpha\,. \tag{1}$$

In this way we obtain a bilinear map called the *hat pairing*:

$$\Omega^p(M) \times \Omega^q(S) \to \Omega^{p+q-k}\big(\mathcal{F}(S,M)\big)\,.$$

An explicit expression of the hat pairing avoiding fiber integration is:

$$(\widehat{\omega\cdot\alpha})_f(Y_f^1,\dots,Y_f^{p+q-k}) = \int_S f^*\big(i_{Y_f^{p+q-k}}\dots i_{Y_f^1}(\omega\circ f)\big)\wedge\alpha\,, \tag{2}$$

for $Y_f^1,\dots Y_f^{p+q-k}$ vector fields on M along $f\in\mathcal{F}(S,M)$. Here we denote by $f^*\beta_f$ the "restricted pull-back" by f of a section β_f of $f^*(\Lambda^m T^*M)$, which is a differential m-form on S given by $f^*\beta_f\colon x\in S\mapsto(\Lambda^m T_x^* f)(\beta_f(x))\in\Lambda^m T_x^* S$, where $T_x^* f\colon T^*_{f(x)}M\to T_x^*S$ denotes the dual of $T_x f$.

The fact that (1) and (2) provide the same differential form on $\mathcal{F}(S,M)$ can be deduced from the identity

$$(\mathrm{ev}^*\,\omega)_{(x,f)}(Y_f^1,\dots,Y_f^{p-k},X_x^1,\dots,X_x^k) = f^*(i_{Y_f^{p-k}}\dots i_{Y_f^1}(\omega\circ f))(X_x^1,\dots,X_x^k)$$

for $Y_f^1,\dots,Y_f^{p-k}\in T_f\mathcal{F}(S,M)$ and $X_x^1,\dots,X_x^k\in T_xS$.

Since $\mathfrak{X}(M)$ acts infinitesimally transitive on the open subset $\mathrm{Emb}(S,M)\subset\mathcal{F}(S,M)$ of embeddings of the k-dimensional oriented manifold S into M, we express $\widehat{\omega}$ at $f\in\mathrm{Emb}(S,M)$ as:

$$(\widehat{\omega\cdot\alpha})_f(X_1\circ f,\dots,X_{p+q-k}\circ f) = \int_S f^*(i_{X_{p+q-k}}\dots i_{X_1}\omega)\wedge\alpha\,. \tag{3}$$

One uses the fact that the "restricted pull-back" by f of $i_{X_{p+q-k}\circ f}\dots i_{X_1\circ f}(\omega\circ f)$ is $f^*(i_{X_{p+q-k}}\dots i_{X_1}\omega)$.

Next we show that the hat pairing is compatible with the exterior derivative of differential forms.

Theorem 1. *The exterior derivative* $\mathbf{d}$ *is a derivation for the hat pairing, i.e.*

$$\mathbf{d}\,(\widehat{\omega\cdot\alpha}) = (\widehat{\mathbf{d}\,\omega})\cdot\alpha + (-1)^p\widehat{\omega\cdot\mathbf{d}\,\alpha}\,, \tag{4}$$

where $\omega\in\Omega^p(M)$ *and* $\alpha\in\Omega^q(S)$.

Proof. Differentiation and fiber integration along the boundary free manifold S commute, so

$$\begin{aligned}\mathbf{d}\,(\widehat{\omega\cdot\alpha}) &= \mathbf{d}\int_S \mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha = \int_S \mathbf{d}\,(\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha)\\ &= \int_S \mathrm{ev}^*\,\mathbf{d}\,\omega\wedge\mathrm{pr}^*\,\alpha + (-1)^p\int_S \mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\mathbf{d}\,\alpha = (\widehat{\mathbf{d}\,\omega})\cdot\alpha + (-1)^p\widehat{\omega\cdot\mathbf{d}\,\alpha}\end{aligned}$$

for all $\omega\in\Omega^p(M)$ and $\alpha\in\Omega^q(S)$. □

The differential form $\widehat{\omega\cdot\alpha}$ is exact if ω is closed and α exact (or if α is closed and ω exact). In the special case $p+q=k$ these conditions imply that the function $\widehat{\omega\cdot\alpha}$ on $\mathcal{F}(S,M)$ vanishes.

Corollary 2. *The hat pairing induces a bilinear map on de Rham cohomology spaces*

$$H^p(M)\times H^q(S)\to H^{p+q-k}\big(\mathcal{F}(S,M)\big)\,. \tag{5}$$

In particular there is a bilinear map

$$H^p(M) \times H^q(M) \to H^{p+q-k}\big(\operatorname{Diff}(M)\big)\,.$$

Remark 3. The cohomology group $H^q(S)$ is isomorphic to the homology group $H_{k-q}(S)$ by Poincaré duality. With the notation $n = k - q$, the hat pairing (5) becomes

$$H^p(M) \times H_n(S) \to H^{p-n}\big(\mathcal{F}(S,M)\big)\,,$$

and it is induced by the map $(\omega, \sigma) \mapsto \fint_\sigma \operatorname{ev}^* \omega$, for differential p-forms ω on M and n-chains σ on S.

If S is a manifold with boundary, then formula (4) receives an extra term coming from integration over the boundary. Let $i_\partial \colon \partial S \to S$ be the inclusion and $r_\partial \colon \mathcal{F}(S,M) \to \mathcal{F}(\partial S, M)$ the restriction map.

Proposition 4. *The identity*

$$\mathbf{d}\,(\widehat{\omega \cdot \alpha}) = (\widehat{\mathbf{d}\,\omega) \cdot \alpha} + (-1)^p \widehat{\omega \cdot \mathbf{d}\,\alpha} + (-1)^{p+q-k} r_\partial^*(\widehat{\omega \cdot i_\partial^* \alpha}^{\,\partial}) \tag{6}$$

holds for $\omega \in \Omega^p(M)$ and $\alpha \in \Omega^q(S)$, where the upper index ∂ assigned to the hat means the pairing

$$\Omega^p(M) \times \Omega^q(\partial S) \to \Omega^{p+q-k+1}\big(\mathcal{F}(\partial S, M)\big)\,.$$

Proof. For any differential n-form β on $S \times \mathcal{F}(S,M)$, the identity

$$\mathbf{d} \fint_S \beta - \fint_S \mathbf{d}\,\beta = (-1)^{n-k} \fint_{\partial S} (i_\partial \times 1_{\mathcal{F}(S,M)})^* \beta$$

holds because of the identity (19) from the appendix. The obvious formulas

$$\operatorname{pr} \circ (i_\partial \times 1_{\mathcal{F}(S,M)}) = i_\partial \circ \operatorname{pr}_\partial, \quad \operatorname{ev} \circ (i_\partial \times 1_{\mathcal{F}(S,M)}) = \operatorname{ev}_\partial\,,$$

for $\operatorname{ev}_\partial \colon \partial S \times \mathcal{F}(S,M) \to M$ and $\operatorname{pr}_\partial \colon \partial S \times \mathcal{F}(S,M) \to \partial S$, are used to compute

$$\begin{aligned}
\mathbf{d}\,(\widehat{\omega \cdot \alpha}) &= \mathbf{d} \fint_S \operatorname{ev}^* \omega \wedge \operatorname{pr}^* \alpha \\
&= \fint_S \mathbf{d}\,(\operatorname{ev}^* \omega \wedge \operatorname{pr}^* \alpha) + (-1)^{p+q-k} \fint_{\partial S} (i_\partial \times 1_{\mathcal{F}(S,M)})^* (\operatorname{ev}^* \omega \wedge \operatorname{pr}^* \alpha) \\
&= \fint_S \operatorname{ev}^* \mathbf{d}\,\omega \wedge \operatorname{pr}^* \alpha + (-1)^p \fint_S \operatorname{ev}^* \omega \wedge \operatorname{pr}^* \mathbf{d}\,\alpha + (-1)^{p+q-k} \fint_{\partial S} \operatorname{ev}_\partial^* \omega \wedge \operatorname{pr}_\partial^* i_\partial^* \alpha \\
&= (\widehat{\mathbf{d}\,\omega) \cdot \alpha} + (-1)^p \widehat{\omega \cdot \mathbf{d}\,\alpha} + (-1)^{p+q-k} r_\partial{}^* (\widehat{\omega \cdot i_\partial^* \alpha}^{\,\partial})\,,
\end{aligned}$$

thus obtaining the requested identity. □

Left $\operatorname{Diff}(M)$ action. The natural left action of the group of diffeomorphisms $\operatorname{Diff}(M)$ on $\mathcal{F}(S,M)$ is $\varphi \cdot f = \varphi \circ f$. The infinitesimal action of $X \in \mathfrak{X}(M)$ is the vector field $\bar{X}$ on $\mathcal{F}(S,M)$:

$$\bar{X}(f) = X \circ f\,, \quad \forall f \in \mathcal{F}(S,M)\,.$$

We denote by $\bar{\varphi}$ the diffeomorphism of $\mathcal{F}(S,M)$ induced by the action of $\varphi \in \operatorname{Diff}(M)$, so $\bar{\varphi}(f) = \varphi \circ f$ is the push-forward by φ.

Proposition 5. *Given $\omega \in \Omega^p(M)$ and $\alpha \in \Omega^q(S)$, the identity*

$$\bar{\varphi}^* \widehat{\omega \cdot \alpha} = \widehat{(\varphi^* \omega) \cdot \alpha} \tag{7}$$

and its infinitesimal version

$$L_{\bar{X}} \widehat{\omega \cdot \alpha} = \widehat{(L_X \omega) \cdot \alpha} \tag{8}$$

hold for all $\varphi \in \mathrm{Diff}(M)$ and $X \in \mathfrak{X}(M)$.

Proof. Using the expression (1) of the hat pairing and identity (15) from the appendix, we have:

$$\begin{aligned} \bar{\varphi}^* \widehat{\omega \cdot \alpha} = \bar{\varphi}^* \fint_S \mathrm{ev}^* \omega \wedge \mathrm{pr}^* \alpha = \fint_S (1_S \times \bar{\varphi})^* (\mathrm{ev}^* \omega \wedge \mathrm{pr}^* \alpha) \\ = \fint_S \mathrm{ev}^* \varphi^* \omega \wedge \mathrm{pr}^* \alpha = \widehat{(\varphi^* \omega) \cdot \alpha}\,, \end{aligned}$$

since $\mathrm{pr} \circ (1_S \times \bar{\varphi}) = \mathrm{pr}$ and $\mathrm{ev} \circ (1_S \times \bar{\varphi}) = \varphi \circ \mathrm{ev}$. □

A similar result is obtained for any smooth map $\eta \in \mathcal{F}(M_1, M_2)$ and its push-forward $\bar{\eta} \colon \mathcal{F}(S, M_1) \to \mathcal{F}(S, M_2)$, $\bar{\eta}(f) = \eta \circ f$:

$$\bar{\eta}^* \widehat{\omega \cdot \alpha} = \widehat{\eta^* \omega \cdot \alpha}\,,$$

for all $\omega \in \Omega^p(M_2)$ and $\alpha \in \Omega^q(S)$.

Lemma 6. *For all vector fields $X \in \mathfrak{X}(M)$, the identity $i_{\bar{X}} \widehat{\omega \cdot \alpha} = \widehat{(i_X \omega) \cdot \alpha}$ holds.*

Proof. The vector field $0_S \times \bar{X}$ on $S \times \mathcal{F}(S, M)$ is ev-related to the vector field X on M, so

$$\begin{aligned} i_{\bar{X}} \widehat{\omega \cdot \alpha} = i_{\bar{X}} \fint_S \mathrm{ev}^* \omega \wedge \mathrm{pr}^* \alpha = \fint_S i_{0_S \times \bar{X}} (\mathrm{ev}^* \omega \wedge \mathrm{pr}^* \alpha) \\ = \fint_S \mathrm{ev}^* (i_X \omega) \wedge \mathrm{pr}^* \alpha = \widehat{(i_X \omega) \cdot \alpha}\,. \end{aligned}$$

At step two we use formula (18) from the appendix. □

Right $\mathrm{Diff}(S)$ action. The natural right action of the diffeomorphism group $\mathrm{Diff}(S)$ on $\mathcal{F}(S, M)$ can be transformed into a left action by $\psi \cdot f = f \circ \psi^{-1}$. The infinitesimal action of $Z \in \mathfrak{X}(S)$ is the vector field $\hat{Z}$ on $\mathcal{F}(S, M)$:

$$\widehat{Z}(f) = -Tf \circ Z\,, \quad \forall f \in \mathcal{F}(S, M)\,.$$

We denote by $\widehat{\psi}$ the diffeomorphism of $\mathcal{F}(S, M)$ induced by the action of ψ, so $\widehat{\psi}(f) = f \circ \psi^{-1}$ is the pull-back by ψ^{-1}.

Proposition 7. *Given $\omega \in \Omega^p(M)$ and $\alpha \in \Omega^q(S)$, the identity*

$$\widehat{\psi}^* \widehat{\omega \cdot \alpha} = \widehat{\omega \cdot \psi^* \alpha}$$

and its infinitesimal version

$$L_{\widehat{Z}} \widehat{\omega \cdot \alpha} = \widehat{\omega \cdot L_Z \alpha}$$

hold for all orientation preserving $\psi \in \mathrm{Diff}(S)$ and $Z \in \mathfrak{X}(S)$.

Proof. The obvious identities $\mathrm{ev}\circ(1_S\times\widehat{\psi})=\mathrm{ev}\circ(\psi^{-1}\times 1_{\mathcal{F}})$, $\mathrm{pr}\circ(1_S\times\widehat{\psi})=\mathrm{pr}$ and $\mathrm{pr}\circ(\psi\times 1_{\mathcal{F}})=\psi\circ\mathrm{pr}$ are used in the computation

$$\begin{aligned}\widehat{\psi^*}\widehat{\omega\cdot\alpha}&=\widehat{\psi}^*\fint_S \mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha=\fint_S(1_S\times\widehat{\psi})^*(\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha)\\&=\fint_S\left((\psi^{-1}\times 1_{\mathcal{F}})^*\,\mathrm{ev}^*\,\omega\right)\wedge\mathrm{pr}^*\,\alpha=\fint_S\mathrm{ev}^*\,\omega\wedge(\psi\times 1_{\mathcal{F}})^*\,\mathrm{pr}^*\,\alpha\\&=\fint_S\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\psi^*\alpha=\widehat{\omega\cdot\psi^*\alpha}\,,\end{aligned}$$

together with formula (17) from the appendix at step four. □

Lemma 8. *The identity* $i_{\widehat{Z}}\widehat{\omega\cdot\alpha}=(-1)^p\widehat{\omega\cdot i_Z\alpha}$ *holds for all vector fields* $Z\in\mathfrak{X}(S)$, *if* $\omega\in\Omega^p(M)$.

Proof. The infinitesimal version of the first identity in the proof of Proposition 7 is $T\,\mathrm{ev}\,.(0_S\times\widehat{Z})=T\,\mathrm{ev}\,.(-Z\times 0_{\mathcal{F}(S,M)})$, so we compute:

$$\begin{aligned}i_{\widehat{Z}}\widehat{\omega\cdot\alpha}&=i_{\widehat{Z}}\fint_S\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha=\fint_S i_{0_S\times\widehat{Z}}(\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha)\\&=\fint_S(i_{0_S\times\widehat{Z}}\,\mathrm{ev}^*\,\omega)\wedge\mathrm{pr}^*\,\alpha=\fint_S(i_{-Z\times 0_{\mathcal{F}(S,M)}}\,\mathrm{ev}^*\,\omega)\wedge\mathrm{pr}^*\,\alpha\\&=\fint_S i_{-Z\times 0_{\mathcal{F}(S,M)}}(\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\alpha)-\fint_S(-1)^p\,\mathrm{ev}^*\,\omega\wedge i_{-Z\times 0_{\mathcal{F}(S,M)}}\,\mathrm{pr}^*\,\alpha\\&=(-1)^p\fint_S\mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*(i_Z\alpha)=(-1)^p\widehat{\omega\cdot i_Z\alpha}\,.\end{aligned}$$

At step two we use formula (18) from the appendix. □

3. Tilda map and hat map

Let $\mathrm{Gr}_k(M)$ be the non-linear Grassmannian (or differentiable Chow variety) of compact oriented k-dimensional submanifolds of M. It is a Fréchet manifold [10] and the tangent space at $N\in\mathrm{Gr}_k(M)$ can be identified with the space of smooth sections of the normal bundle $TN^\perp=(TM|_N)/TN$. The tangent vector at N determined by the section $Y_N\in\Gamma(TM|_N)$ is denoted by $\tilde{Y}_N\in T_N\,\mathrm{Gr}_k(M)$.

The *tilda map* [6] associates to any p-form ω on M a $(p-k)$-form $\tilde{\omega}$ on $\mathrm{Gr}_k(M)$ by:

$$\tilde{\omega}_N(\tilde{Y}_N^1,\dots,\tilde{Y}_N^{p-k})=\int_N i_{Y_N^{p-k}}\cdots i_{Y_N^1}\omega\,. \tag{9}$$

Here all $\tilde{Y}_N^j$ are tangent vectors at $N\in\mathrm{Gr}_k(M)$, i.e. sections of $TN^\perp$ represented by sections Y_N^j of $TM|_N$. Then $i_{Y_N^{p-k}}\cdots i_{Y_N^1}\omega\in\Omega^k(N)$ does not depend on representatives Y_N^j of $\tilde{Y}_N^j$, and integration is well defined since $N\in\mathrm{Gr}_k(M)$ comes with an orientation.

Let S be a compact oriented k-dimensional manifold. The *hat map* is the hat pairing with the constant function $1\in\Omega^0(S)$. It associates to any form $\omega\in\Omega^p(M)$

the form $\widehat{\omega} \in \Omega^{p-k}(\mathcal{F}(S,M))$:

$$\widehat{\omega} = \widehat{\omega \cdot 1} = \fint_S \mathrm{ev}^*\,\omega\,. \tag{10}$$

On the open subset $\mathrm{Emb}(S,M) \subset \mathcal{F}(S,M)$ of embeddings, formula (2) gives

$$\widehat{\omega}(X_1 \circ f, \ldots, X_{p-k} \circ f) = \int_S f^*(i_{X_{p-k}} \ldots i_{X_1}\omega)\,. \tag{11}$$

Remark 9. The hat map induces a transgression on cohomology spaces

$$H^p(M) \to H^{p-k} = \big(\mathcal{F}(S,M)\big)\,.$$

When S is the circle, then one obtains the usual transgression map with values in the $(p-1)$-th cohomology space of the free loop space of M.

Let π denote the natural map

$$\pi\colon\ \mathrm{Emb}(S,M) \to \mathrm{Gr}_k(M), \quad \pi(f) = f(S)\,.$$

where the orientation on $f(S)$ is chosen such that the diffeomorphism $f\colon S \to f(S)$ is orientation preserving. The image $\pi(\mathrm{Emb}(S,M))$ is the manifold $\mathrm{Gr}_k^S(M)$ of k-dimensional submanifolds of M of type S. Then $\pi\colon\ \mathrm{Emb}(S,M) \to \mathrm{Gr}_k^S(M)$ is a principal bundle over $\mathrm{Gr}_k^S(M)$ with structure group $\mathrm{Diff}_+(S)$, the group of orientation preserving diffeomorphisms of S.

Note that there is a natural action of the group $\mathrm{Diff}(M)$ on the non-linear Grassmannian $\mathrm{Gr}_k(M)$ given by $\varphi \cdot N = \varphi(N)$. Let $\tilde{\varphi}$ be the diffeomorphism of $\mathrm{Gr}_k(M)$ induced by the action of $\varphi \in \mathrm{Diff}(M)$. Then $\tilde{\varphi} \circ \pi = \pi \circ \bar{\varphi}$ for the restriction of $\bar{\varphi}(f) = \varphi \circ f$ to a diffeomorphism of $\mathrm{Emb}(S,M) \subset \mathcal{F}(S,M)$. As a consequence, the infinitesimal generators for the $\mathrm{Diff}(M)$ actions on $\mathrm{Gr}_k(M)$ and on $\mathrm{Emb}(S,M)$ are π-related. This means that for all $X \in \mathfrak{X}(M)$, the vector fields $\tilde{X}$ on $\mathrm{Gr}_k(M)$ given by $\tilde{X}(N) = X|_N$ and $\bar{X}$ on $\mathrm{Emb}(S,M)$ given by $\bar{X}(f) = X \circ f$ are π-related.

Proposition 10. *The hat map on* $\mathrm{Emb}(S,M)$ *and the tilda map on* $\mathrm{Gr}_k(M)$ *are related by* $\widehat{\omega} = \pi^*\tilde{\omega}$*, for any* k*-dimensional oriented manifold* S*.*

Proof. For the proof we use the fact that $\mathfrak{X}(M)$ acts infinitesimally transitive on $\mathrm{Emb}(S,M)$, so $T_f\,\mathrm{Emb}(S,M) = \{X \circ f\colon X \in \mathfrak{X}(M)\}$. With (9) and (11) we compute:

$$\begin{aligned}(\pi^*\tilde{\omega})_f(X_1 \circ f, \ldots, X_{p-k} \circ f) &= \tilde{\omega}_{f(S)}(X_1|_{f(S)}, \ldots, X_{p-k}|_{f(S)}) \\ &= \int_{f(S)} i_{X_{p-k}} \ldots i_{X_1}\omega = \int_S f^*(i_{X_{p-k}} \ldots i_{X_1}\omega) = \widehat{\omega}_f(X_1 \circ f, \ldots, X_{p-k} \circ f)\,,\end{aligned}$$

since $\bar{X}$ and $\tilde{X}$ are π-related. □

From the properties of the hat pairing presented in Proposition 5, Lemma 6 and Theorem 1, a hat calculus follows easily:

Proposition 11. *For any* $\omega \in \Omega^p(M)$*,* $\varphi \in \mathrm{Diff}(M)$*,* $X \in \mathfrak{X}(M)$*, and* $\eta \in \mathcal{F}(M',M)$ *with push-forward* $\bar{\eta}\colon \mathcal{F}(S,M') \to \mathcal{F}(S,M)$*, the following identities hold:*

(1) $\bar{\varphi}^*\widehat{\omega} = \widehat{\varphi^*\omega}$ *and* $\bar{\eta}^*\widehat{\omega} = \widehat{\eta^*\omega}$

(2) $L_{\bar{X}}\widehat{\omega} = \widehat{L_X\omega}$

(3) $i_{\bar{X}}\widehat{\omega} = \widehat{i_X\omega}$

(4) $\mathbf{d}\,\widehat{\omega} = \widehat{\mathbf{d}\,\omega}$.

Remark 12. If S is a manifold with boundary, then the formula 4. above receives an extra term coming from integration over the boundary ∂S as in Proposition 4:

$$\mathbf{d}\,\widehat{\omega} = \widehat{\mathbf{d}\,\omega} + (-1)^{p-k} r_\partial^* \widehat{\omega}^\partial \tag{12}$$

for $\omega \in \Omega^p(M)$. As before, $r_\partial \colon \mathcal{F}(S,M) \to \mathcal{F}(\partial S, M)$ denotes the restriction map on functions and $\omega \in \Omega^p(M) \mapsto \widehat{\omega}^\partial \in \Omega^{p-k+1}(\mathcal{F}(\partial S, M))$.

Now the properties of the tilda calculus follow imediately from Proposition 11.

Proposition 13. [6] *For any* $\omega \in \Omega^p(M)$, $\varphi \in \mathrm{Diff}(M)$ *and* $X \in \mathfrak{X}(M)$, *the following identities hold:*

(1) $\tilde{\varphi}^*\tilde{\omega} = \widetilde{\varphi^*\omega}$

(2) $L_{\tilde{X}}\tilde{\omega} = \widetilde{L_X\omega}$

(3) $i_{\tilde{X}}\tilde{\omega} = \widetilde{i_X\omega}$

(4) $\mathbf{d}\,\tilde{\omega} = \widetilde{\mathbf{d}\,\omega}$.

Proof. We verify the identities 1. and 4. From relation 1. from Proposition 11 we get that

$$\pi^*\tilde{\varphi}^*\tilde{\omega} = \bar{\varphi}^*\pi^*\tilde{\omega} = \bar{\varphi}^*\widehat{\omega} = \widehat{\varphi^*\omega} = \pi^*\widetilde{\varphi^*\omega}\,,$$

and this implies the first identity. Using identity 4. from Proposition 11 we compute

$$\pi^*\mathbf{d}\,\tilde{\omega} = \mathbf{d}\,\pi^*\tilde{\omega} = \mathbf{d}\,\widehat{\omega} = \widehat{\mathbf{d}\,\omega} = \pi^*\widetilde{\mathbf{d}\,\omega}\,,$$

which shows the last identity. □

Hamiltonian formalism for p-branes. In this section we show how the hat calculus appears in the hamiltonian formalism for p-branes and open p-branes [1] [2].

Let S be a compact oriented p-dimensional manifold. The phase space for the p-brane world volume $S \times \mathbb{R}$ is the cotangent bundle $T^*\mathcal{F}(S,M)$, where the canonical symplectic form is twisted. The twisting consists in adding a magnetic term, namely the pull-back of a closed 2-form on the base manifold, to the canonical symplectic form on a cotangent bundle [11]. These twisted symplectic forms appear also in cotangent bundle reduction.

We consider a closed differential form $H \in \Omega^{p+2}(M)$. Since $\dim S = p$, the hat map (10) provides a closed 2-form $\widehat{H}$ on $\mathcal{F}(S,M)$. If $\pi_{\mathcal{F}} : T^*\mathcal{F}(S,M) \to \mathcal{F}(S,M)$ denotes the canonical projection, the twisted symplectic form on $T^*\mathcal{F}(S,M)$ is

$$\Omega_H = -\mathbf{d}\,\Theta_{\mathcal{F}} + \frac{1}{2}\pi_{\mathcal{F}}^*\widehat{H}\,,$$

where $\Theta_{\mathcal{F}}$ is the canonical 1-form on $T^*\mathcal{F}(S,M)$.

For the description of open branes one considers a compact oriented p-dimensional manifold S with boundary ∂S and a submanifold D of M. The phase space is in this case the cotangent bundle $T^*\mathcal{F}_D(S,M)$ over the manifold [13]

$$\mathcal{F}_D(S,M) = \{f : S \to M | f(\partial S) \subset D\}.$$

The twisting of the canonical symplectic form is done with a closed differential form $H \in \Omega^{p+2}(M)$ with $i^*H = \mathbf{d}\,B$ for some $B \in \Omega^{p+1}(D)$, where $i\colon D \to M$ denotes the inclusion. The twisted symplectic form on $T^*\mathcal{F}_D(S,M)$ is

$$\Omega_{(H,B)} = -\mathbf{d}\,\Theta_{\mathcal{F}_D} + \frac{1}{2}\pi^*_{\mathcal{F}_D}(\widehat{H} - \partial^*\widehat{B}^{\partial})$$

with $\partial\colon \mathcal{F}_D(S,M) \to \mathcal{F}(\partial S, D)$ the restriction map and $\pi_{\mathcal{F}_D}\colon T^*\mathcal{F}_D(S,M) \to \mathcal{F}_D(S,M)$. To distinguish between the hat calculus for $\mathcal{F}(S,M)$ and the hat calculus for $\mathcal{F}(\partial S, M)$, we denote $\widehat{\ }^{\partial}\colon \Omega^n(M) \to \Omega^{n-p+1}(\mathcal{F}(\partial S, M))$.

The only thing we have to verify is the closedness of $\widehat{H} - \partial^*\widehat{B}^{\partial}$. We first notice that (12) implies $\mathbf{d}\,\widehat{H} = \widehat{\mathbf{d}\,H} + r^*_{\partial}\widehat{H}^{\partial}$, where $r_{\partial}\colon \mathcal{F}(S,M) \to \mathcal{F}(\partial S, M)$ denotes the restriction map, and identity 4 from Proposition 11 implies $\widehat{\mathbf{d}\,B}^{\partial} = \mathbf{d}\,\widehat{B}^{\partial}$. On the other hand identity 1 from Proposition 11 ensures that $\widehat{i^*H}^{\partial} = \bar{i}^*\widehat{H}^{\partial}$, with $\bar{i}\colon \mathcal{F}(\partial S, D) \to \mathcal{F}(\partial S, M)$ denoting the push-forward by $i\colon D \to M$. Knowing that $r_{\partial} = \bar{i} \circ \partial$, we compute:

$$\mathbf{d}\,\widehat{H} = \widehat{\mathbf{d}\,H} + r^*_{\partial}\widehat{H}^{\partial} = \partial^*\bar{i}^*\widehat{H}^{\partial} = \partial^*\widehat{i^*H}^{\partial} = \partial^*\widehat{\mathbf{d}\,B}^{\partial} = \mathbf{d}\,\partial^*\widehat{B}^{\partial}\,,$$

so the closed 2-form $\widehat{H} - \partial^*\widehat{B}^{\partial}$ provides a twist for the canonical symplectic form on the cotangent bundle $T^*\mathcal{F}_D(S,M)$.

Non-linear Grassmannians as symplectic manifolds. In this subsection we recall properties of the co-dimension two non-linear Grassmannian as a symplectic manifold.

Proposition 14 ([8])**.** *Let M be a closed m-dimensional manifold with volume form ν. The tilda map provides a symplectic form $\tilde{\nu}$ on* $\mathrm{Gr}_{m-2}(M)$

$$\tilde{\nu}_N(\tilde{X}_N, \tilde{Y}_N) = \int_N i_{Y_N} i_{X_N} \nu\,,$$

for $\tilde{X}_N$ and $\tilde{Y}_N$ sections of $TN^{\perp}$ determined by sections X_N and Y_N of $TM|_N$.

Proof. The 2-form $\tilde{\nu}$ is closed since $\mathbf{d}\,\tilde{\nu} = \widetilde{\mathbf{d}\,\nu}$ by the tilda calculus. To verify that it is also (weakly) non-degenerate, let X_N be an arbitrary vector field along N such that $\int_N i_{Y_N} i_{X_N}\nu = 0$ for all vector fields Y_N along N. Then X_N must be tangent to N, so $\tilde{X}_N = 0$. □

In dimension $m = 3$ the symplectic form $\tilde{\nu}$ is known as the Marsden–Weinstein symplectic from on the space of unparameterized oriented links, see [12], [3].

Hamiltonian $\mathrm{Diff}_{\mathrm{ex}}(M,\nu)$ action. The action of the group $\mathrm{Diff}(M,\nu)$ of volume preserving diffeomorphisms of M on $\mathrm{Gr}_{m-2}(M)$ preserves the symplectic form $\tilde{\nu}$:

$$\tilde{\varphi}^*\tilde{\nu} = \widetilde{\varphi^*\nu} = \tilde{\nu}\,, \quad \forall \varphi \in \mathrm{Diff}(M,\nu)\,.$$

The subgroup $\mathrm{Diff}_{\mathrm{ex}}(M,\nu)$ of exact volume preserving diffeomorphisms acts in a hamiltonian way on the symplectic manifold $(\mathrm{Gr}_{m-2}(M),\tilde{\nu})$. Its Lie algebra is $\mathfrak{X}_{\mathrm{ex}}(M,\nu)$, the Lie algebra of exact divergence free vector fields, i.e. vector fields X_α such that $i_{X_\alpha}\nu = \mathbf{d}\,\alpha$ for a potential form $\alpha \in \Omega^{m-2}(M)$. The infinitesimal action of X_α is the vector field $\tilde{X}_\alpha$. By the tilda calculus $\tilde{\alpha} \in \mathcal{F}(\mathrm{Gr}_{m-2}(M))$ is a hamiltonian function for the hamiltonian vector field $\tilde{X}_\alpha$:

$$i_{\tilde{X}_\alpha}\tilde{\nu} = \widetilde{i_{X_\alpha}\nu} = \widetilde{\mathbf{d}\,\alpha} = \mathbf{d}\,\tilde{\alpha}\,.$$

It depends on the particular choice of the potential α of X_α. A fixed continuous right inverse $b : \mathbf{d}\,\Omega^{m-2}(M) \to \Omega^{m-2}(M)$ to the differential $\mathbf{d}$ picks up a potential $b(\mathbf{d}\,\alpha)$ of X_α. The corresponding momentum map is:

$$\mathbf{J} : \mathcal{M} \to \mathfrak{X}_{\mathrm{ex}}(M,\nu)^*\,, \quad \langle \mathbf{J}(N), X_\alpha\rangle = \widetilde{b(\mathbf{d}\,\alpha)}(N) = \int_N b(\mathbf{d}\,\alpha)\,.$$

On the connected component $\mathcal{M}$ of $N \in \mathrm{Gr}_{m-2}(M)$, the non-equivariance of $\mathbf{J}$ is measured by the Lie algebra 2-cocycle on $\mathfrak{X}_{\mathrm{ex}}(M,\nu)$

$$\begin{aligned}\sigma_N(X,Y) &= \langle \mathbf{J}(N), [X,Y]^{\mathrm{op}}\rangle - \tilde{\nu}(\tilde{X},\tilde{Y})(N) = (\widetilde{b\,\mathbf{d}\,i_Y i_X \nu})(N) - (\widetilde{i_Y i_X \nu})(N)\\ &= (\widetilde{P i_X i_Y \nu})(N) = \int_N P i_X i_Y \nu\,.\end{aligned}$$

Here $P = 1_{\Omega^{m-2}(M)} - b \circ \mathbf{d}$ is a continuous linear projection on the subspace of closed $(m-2)$-forms and $(X,Y) \mapsto [P i_Y i_X \nu] \in H^{m-2}(M)$ is the universal Lie algebra 2-cocycle on $\mathfrak{X}_{\mathrm{ex}}(M,\nu)$ [14]. The cocycle σ_N is cohomologous to the Lichnerowicz cocycle

$$\sigma_\eta(X,Y) = \int_M \eta(X,Y)\nu\,, \tag{13}$$

where η is a closed 2-form Poincaré dual to N [15].

If ν is an integral volume form, then σ_N is integrable [8]. The connected component $\mathcal{M}$ of $\mathrm{Gr}_{m-2}(M)$ is a coadjoint orbit of a 1-dimensional central Lie group extension of $\mathrm{Diff}_{\mathrm{ex}}(M,\nu)$ integrating σ_N, and $\tilde{\nu}$ is the Kostant-Kirillov-Souriau symplectic form. [6].

4. Bar map

When a volume form μ on the compact k-dimensional manifold S is given, one can associate to each differential p-form on M a differential p-form on $\mathcal{F}(S,M)$

$$\bar{\omega}(Y_f^1,\dots,Y_f^p) = \int_S \omega(Y_f^1,\dots,Y_f^p)\mu\,, \quad \forall Y_f^i \in T_f\mathcal{F}(S,M)\,,$$

where $\omega(Y_f^1,\dots,Y_f^p)\colon x\mapsto \omega_{f(x)}(Y_f^1(x),\dots,Y_f^p(x))$ defines a smooth function on S. In this way a *bar map* is defined. Formula (2) assures that this bar map is just the hat pairing of differential forms on M with the volume form μ

$$\bar\omega = \widehat{\omega\cdot\mu} = \int_S \mathrm{ev}^*\,\omega\wedge\mathrm{pr}^*\,\mu\,. \tag{14}$$

From the properties of the hat pairing presented in Proposition 5, Lemma 6 and Theorem 1, one can develop a bar calculus.

Proposition 15. *For any $\omega\in\Omega^p(M)$, $\varphi\in\mathrm{Diff}(M)$ and $X\in\mathfrak{X}(M)$, the following identities hold:*

(1) $\bar\varphi^*\bar\omega = \overline{\varphi^*\omega}$
(2) $L_{\bar X}\bar\omega = \overline{L_X\omega}$
(3) $i_{\bar X}\bar\omega = \overline{i_X\omega}$
(4) $\mathbf{d}\,\bar\omega = \overline{\mathbf{d}\,\omega}$.

$\mathcal{F}(S,M)$ as symplectic manifold. Let (M,ω) be a connected symplectic manifold and S a compact k-dimensional manifold with a fixed volume form μ, normalized such that $\int_S\mu=1$. The following fact is well known:

Proposition 16. *The bar map provides a symplectic form $\bar\omega$ on $\mathcal{F}(S,M)$:*

$$\bar\omega_f(X_f,Y_f) = \int_S \omega(X_f,Y_f)\mu\,.$$

Proof. That $\bar\omega$ is closed follows from the bar calculus: $\mathbf{d}\,\bar\omega = \overline{\mathbf{d}\,\omega}=0$. The (weakly) non-degeneracy of $\bar\omega$ can be verified as follows. If the vector field X_f on M along S is non-zero, then $X_f(x)\neq 0$ for some $x\in S$. Because ω is non-degenerate, one can find another vector field Y_f along f such that $\omega(X_f,Y_f)$ is a bump function on S. Then $\bar\omega(X_f,Y_f)=\int_S\omega(X_f,Y_f)\mu\neq 0$, so X_f does not belong to the kernel of $\bar\omega$, thus showing that the kernel of $\bar\omega$ is trivial. □

Hamiltonian action on M. Let G be a Lie group acting in a hamiltonian way on M with momentum map $J\colon M\to\mathfrak{g}^*$. Then $\mathcal{F}(S,M)$ inherits a G-action: $(g\cdot f)(x)=g\cdot(f(x))$ for any $x\in S$. The infinitesimal generator is $\xi_{\mathcal{F}}=\bar\xi_M$ for any $\xi\in\mathfrak{g}$, where ξ_M denotes the infinitesimal generator for the G-action on M. The bar calculus shows quickly that G acts in a hamiltonian way on $\mathcal{F}(S,M)$ with momentum map

$$\mathbf{J}=\bar J\colon \mathcal{F}(S,M)\to\mathfrak{g}^*\,,\quad \bar J(f)=\int_S (J\circ f)\mu\,,\quad \forall f\in\mathcal{F}(S,M)\,.$$

Indeed, for all $\xi\in\mathfrak{g}$

$$i_{\xi_{\mathcal{F}}}\bar\omega = i_{\bar\xi_M}\bar\omega = \overline{i_{\xi_M}\omega} = \overline{\mathbf{d}\,\langle J,\xi\rangle} = \mathbf{d}\,\langle\bar J,\xi\rangle\,.$$

Let M be connected and let σ be the $\mathbb{R}$-valued Lie algebra 2-cocycle on $\mathfrak{g}$ measuring the non-equivariance of J, i.e.

$$\sigma(\xi,\eta)=\langle J(x),[\xi,\eta]\rangle-\omega(\xi_M,\eta_M)(x)\,,\quad x\in M\,,$$

(both terms are hamiltonian function for the vector field $[\xi,\eta]_M = -[\xi_M,\eta_M]$). Then the non-equivariance of $\mathbf{J} = \bar{J}$ is also measured by σ: for all $f \in \mathcal{F}(S,M)$

$$\langle \bar{J}(f), [\xi,\eta]\rangle - \bar{\omega}(\xi_{\mathcal{F}},\eta_{\mathcal{F}})(f) = \overline{\langle J,[\xi,\eta]\rangle}(f) - \overline{\omega(\xi_M,\eta_M)}(f) = \sigma(\xi,\eta)\,.$$

Hamiltonian $\mathrm{Diff}_{\mathrm{ham}}(M,\omega)$ action. The action of the group $\mathrm{Diff}(M,\omega)$ of symplectic diffeomorphisms preserves the symplectic form $\bar{\omega}$:

$$\bar{\varphi}^*\bar{\omega} = \overline{\varphi^*\omega} = \bar{\omega}\,, \quad \forall \varphi \in \mathrm{Diff}(M,\omega)\,.$$

The subgroup $\mathrm{Diff}_{\mathrm{ham}}(M,\omega)$ of hamiltonian diffeomorphisms of M acts in a hamiltonian way on the symplectic manifold $\mathcal{F}(S,M)$. The infinitesimal action of $X_h \in \mathfrak{X}_{\mathrm{ham}}(M,\omega)$, $h \in \mathcal{F}(M)$, is the hamiltonian vector field $\bar{X}_h$ on $\mathcal{F}(S,M)$ with hamiltonian function $\bar{h}$. This follows by the bar calculus:

$$\mathbf{d}\,\bar{h} = \overline{\mathbf{d}\,h} = \overline{i_{X_h}\omega} = i_{\bar{X}_h}\bar{\omega}\,.$$

The hamiltonian function $\bar{h}$ of $\bar{X}_h$ depends on the particular choice of the hamiltonian function h. To solve this problem we fix a point $x_0 \in M$ and we choose the unique hamiltonian function h with $h(x_0) = 0$, since M is connected. The corresponding momentum map is

$$\mathbf{J} : \mathcal{F}(S,M) \to \mathfrak{X}_{\mathrm{ham}}(M,\omega)^*\,, \quad \langle \mathbf{J}(f), X_h\rangle = \bar{h}(f) = \int_S (h\circ f)\mu\,.$$

The Lie algebra 2-cocycle on $\mathfrak{X}_{\mathrm{ham}}(M,\omega)$ measuring the non-equivariance of the momentum map is

$$\sigma(X,Y) = -\omega(X,Y)(x_0)\,,$$

by the bar calculus

$$\begin{aligned}\sigma(X,Y)(f) &= \langle \mathbf{J}(f), [X,Y]^{\mathrm{op}}\rangle - \bar{\omega}(X_{\mathcal{F}},Y_{\mathcal{F}})(f)\\ &= \overline{\omega(X,Y) - \omega(X,Y)(x_0)}(f) - \bar{\omega}(\bar{X},\bar{Y})(f) = -\omega(X,Y)(x_0)\,.\end{aligned}$$

This is a Lie algebra cocycle describing the central extension

$$0 \to \mathbb{R} \to \mathcal{F}(M) \to \mathfrak{X}_{\mathrm{ham}}(M,\omega) \to 0$$

where $\mathcal{F}(M)$ is enowed with the canonical Poisson bracket. A group cocycle on $\mathrm{Diff}_{\mathrm{ham}}(M,\omega)$ integrating the Lie algebra cocycle σ if ω exact is studied in [9].

Hamiltonian $\mathrm{Diff}_{\mathrm{ex}}(S,\mu)$ action. The (left) action of the group $\mathrm{Diff}(S,\mu)$ of volume preserving diffeomorphisms preserves the symplectic form $\bar{\omega}$:

$$\widehat{\psi}^*\bar{\omega} = \widehat{\psi}^*\widehat{\omega\cdot\mu} = \widehat{\omega\cdot\psi^*\mu} = \widehat{\omega\cdot\mu} = \bar{\omega}\,, \quad \forall \psi \in \mathrm{Diff}(S,\mu)\,.$$

The subgroup $\mathrm{Diff}_{\mathrm{ex}}(S,\mu)$ of exact volume preserving diffeomorphisms acts in a hamiltonian way on the symplectic manifold $\mathcal{F}(S,M)$. The infinitesimal action of the exact divergence free vector field $X_\alpha \in \mathfrak{X}_{\mathrm{ex}}(S,\mu)$ with potential form $\alpha \in \Omega^{k-2}(S)$ is the hamiltonian vector field $\widehat{X}_\alpha$ on $\mathcal{F}(S,M)$ with hamiltonian function $\widehat{\omega\cdot\alpha}$. Indeed, from $i_{X_\alpha}\mu = \mathbf{d}\,\alpha$ follows by the hat calculus that

$$\mathbf{d}\,(\widehat{\omega\cdot\alpha}) = \widehat{\mathbf{d}\,\omega\cdot\alpha} + \widehat{\omega\cdot\mathbf{d}\,\alpha} = \omega\cdot\widehat{i_{X_\alpha}\mu} = i_{\widehat{X}_\alpha}\widehat{\omega\cdot\mu} = i_{\widehat{X}_\alpha}\bar{\omega}\,.$$

If the symplectic form ω is exact, then the corresponding momentum map is

$$\mathbf{J}\colon \mathcal{F}(S,M) \to \mathfrak{X}_{\mathrm{ex}}(S,\mu)^*, \quad \langle \mathbf{J}(f), X_\alpha\rangle = \widehat{(\omega\cdot\alpha)}(f) = \int_S f^*\omega\wedge\alpha.$$

It takes values in the regular part of $\mathfrak{X}_{\mathrm{ex}}(S,\mu)^*$, which can be identified with $\mathbf{d}\,\Omega^1(S)$, so we can write $\mathbf{J}(f) = f^*\omega$ under this identification.

In general the hamiltonian function $\widehat{\omega\cdot\alpha}$ of $\widehat{X}_\alpha$ depends on the particular choice of the potential form α of X_α. To fix this problem we consider as in Section 3 a continuous right inverse $b : \mathbf{d}\,\Omega^{m-2}(M) \to \Omega^{m-2}(M)$ to the differential $\mathbf{d}$, so $b(\mathbf{d}\,\alpha)$ is a potential for X_α. The corresponding momentum map is

$$\mathbf{J} : \mathcal{F}(S,M) \to \mathfrak{X}_{\mathrm{ex}}(S,\mu)^*, \quad \langle \mathbf{J}(f), X_\alpha\rangle = (\widehat{\omega\cdot b\mathbf{d}\,\alpha})(f) = \int_S f^*\omega\wedge b(\mathbf{d}\,\alpha)\,.$$

On a connected component $\mathcal{F}$ of $\mathcal{F}(S,M)$, the non-equivariance of $\mathbf{J}$ is measured by the Lie algebra 2-cocycle

$$\begin{aligned}\sigma_{\mathcal{F}}(X,Y) &= \langle \mathbf{J}(f), [X,Y]\rangle - \bar{\omega}(\hat{X},\hat{Y})(f) = (\omega\cdot b\mathbf{d}\,i_Y i_X\mu)\hat{}(f) - (\omega\cdot i_Y i_X\mu)\hat{}(f)\\ &= (\omega\cdot Pi_X i_Y\mu)\hat{}(f) = \int_S f^*\omega\wedge Pi_X i_Y\mu\end{aligned}$$

on the Lie algebra of exact divergence free vector fields, for $P = 1 - b\mathbf{d}$ the projection on the subspace of closed $(m-2)$-forms. It does not depend on $f\in\mathcal{F}$, because the cohomology class $[f^*\omega]\in H^2(S)$ does not depend on the choice of f. The cocycle $\sigma_{\mathcal{F}}$ is cohomologous to the Lichnerowicz cocycle $\sigma_{f^*\omega}$ defined in (13) [15]. Since $\int_S\mu = 1$, the cocycle $\sigma_{\mathcal{F}}$ is integrable if and only if the cohomology class of $f^*\omega$ is integral [8].

Remark 17. The two equivariant momentum maps on the symplectic manifold $\mathcal{F}(S,M)$, for suitable central extensions of the hamiltonian group $\mathrm{Diff}_{\mathrm{ham}}(M,\omega)$ and of the group $\mathrm{Diff}_{\mathrm{ex}}(S,\mu)$ of exact volume preserving diffeomorphisms, form the dual pair for ideal incompressible fluid flow [12] [4].

5. Appendix: Fiber integration

Chapter VII in [5] is devoted to the concept of integration over the fiber in locally trivial bundles. We particularize this fiber integration to the case of trivial bundles $S\times M\to M$, listing its main properties without proofs.

Let S be a compact k-dimensional manifold. Fiber integration over S assigns to $\omega\in\Omega^n(S\times M)$ the differential form $\int\!\!\!\!\!\!\int_S\omega\in\Omega^{n-k}(M)$ defined by

$$(\int\!\!\!\!\!\!\int_S\omega)(x) = \int_S \omega_x \in \Lambda^{n-k}T_x^*M\,,\quad \forall x\in M\,,$$

where $\omega_x\in\Omega^k(S,\Lambda^{n-k}T_x^*M)$ is the retrenchment of ω to the fiber over x:

$$\langle\omega_x(Z_s^1,\dots,Z_s^{n-k}), X_x^1\wedge\cdots\wedge X_x^k\rangle = \omega_{(s,x)}(X_x^1,\dots,X_x^k,Z_s^1,\dots,Z_s^{n-k})$$

for all $X_x^i\in T_xM$ and $Z_s^j\in T_sS$.

The properties of the fiber integration used in the text are special cases of the Propositions (VIII) and (X) in [5]:

- Pull-back of fiber integrals:

$$f^* \fint_S \omega = \fint_S (1_S \times f)^* \omega \,, \quad \forall f \in \mathcal{F}(M', M) \,, \tag{15}$$

with infinitesimal version

$$L_X \fint_S \omega = \fint_S L_{0_S \times X} \omega \,, \quad \forall X \in \mathfrak{X}(M) \,. \tag{16}$$

- Invariance under pull-back by orientation preserving diffeomorphisms of S:

$$\fint_S (\varphi \times 1_M)^* \omega = \fint_S \omega \,, \quad \forall \varphi \in \mathrm{Diff}_+(S) \,, \tag{17}$$

with infinitesimal version $\fint_S L_{Z \times 0_M} \omega = 0 \,, \quad \forall Z \in \mathfrak{X}(S)$.

- Insertion of vector fields into fiber integrals:

$$i_X \fint_S \omega = \fint_S i_{0_S \times X} \omega, \quad \forall X \in \mathfrak{X}(M). \tag{18}$$

- Integration along boundary free manifolds commutes with differentiation. When ∂S denotes the boundary of the k-dimensional compact manifold S and $i_\partial \colon \partial S \to S$ the inclusion,

$$\mathbf{d} \fint_S \beta - \fint_S \mathbf{d}\, \beta = (-1)^{n-k} \fint_{\partial S} (i_\partial \times 1_M)^* \beta \tag{19}$$

holds for any differential n-form β on $S \times M$.

References

[1] Alekseev, A., Strobl, T., *Current algebras and differential geometry*, J. High Energy Phys. **03** (2005), 14.

[2] Bonelli, G., Zabzine, M., *From current algebras for p–branes to topological M–theory*, J. High Energy Phys. **09** (2005), 15.

[3] Brylinski, J.-L., *Loop spaces, characteristic classes and geometric quantization*, vol. 107, Progr. Math., Birkhäuser, 1993.

[4] Gay-Balmaz, F., Vizman, C., *Dual pairs in fluid dynamics*, Ann. Global Anal. Geom. (2011), DOI:10.1007/s10455-011-9267-z.

[5] Greub, W., Halperin, S., Vanstone, R., *Connections, curvature, and cohomology. Vol. I: De Rham cohomology of manifolds and vector bundles*, Pure and Applied Mathematics, vol. 47, Academic Press, New York–London, 1972.

[6] Haller, S., Vizman, C., *Non-linear Grassmannians as coadjoint orbits*, Math. Ann. **329** (2004), 771–785.

[7] Hirsch, M. W., *Differential topology*, vol. 33, Springer, 1976, Grad. Texts in Math.

[8] Ismagilov, R. S., *Representations of infinite–dimensional groups*, vol. 152, Ams. Math. Soc., 1996, Translations of Mathematical Monographs.

[9] Ismagilov, R. S., Losik, M., Michor, P. W., *A 2–cocycle on a group of symplectomorphisms*, Moscow Math. J. **6** (2006), 307–315.

[10] Kriegl, A., Michor, P. W., *The convenient setting of global analysis*, vol. 53, Math. Surveys Monogr., 1997.

[11] Marsden, J. E., Ratiu, T., *Introduction to Mechanics and Symmetry*, 2nd ed., Springer, 1999.

[12] Marsden, J. E., Weinstein, A., *Coadjoint orbits, vortices, and Clebsch variables for incompressible fluids*, Phys. D **7** (1983), 305–323.

[13] Michor, P. W., *Manifolds of differentiable mappings*, vol. 3, Shiva Math. Series, Shiva Publishing Ltd., Nantwich, 1980.

[14] Roger, C., *Extensions centrales d'algèbres et de groupes de Lie de dimension infinie, algèbre de Virasoro et généralisations*, Rep. Math. Phys. **35** (1995), 225–266.

[15] Vizman, C., *Lichnerowicz cocycles and central Lie group extensions*, An. Univ. Vest Timis., Ser. Mat.–Inform. **48** (1–2) (2010), 285–297.

West University of Timişoara, Department of Mathematics
E-mail: vizman@math.uvt.ro