ARCHIVUM MATHEMATICUM (BRNO)
Tomus 48 (2012), 113–120

AN ELEMENTARY PROOF OF A CONGRUENCE BY SKULA AND GRANVILLE

Romeo Meštrović

Abstract. Let $p \geq 5$ be a prime, and let $q_p(2) := (2^{p-1}-1)/p$ be the Fermat quotient of p to base 2. The following curious congruence was conjectured by L. Skula and proved by A. Granville

$$q_p(2)^2 \equiv -\sum_{k=1}^{p-1} \frac{2^k}{k^2} \pmod{p}.$$

In this note we establish the above congruence by entirely elementary number theory arguments.

1. Introduction and statement of the main result

The Fermat Little Theorem states that if p is a prime and a is an integer not divisible by p, then $a^{p-1} \equiv 1 \pmod{p}$. This gives rise to the definition of the *Fermat quotient* of p to base a

$$q_p(a) := \frac{a^{p-1}-1}{p},$$

which is an integer. Fermat quotients played an important role in the study of cyclotomic fields and Fermat Last Theorem. More precisely, divisibility of Fermat quotient $q_p(a)$ by p has numerous applications which include the Fermat Last Theorem and squarefreeness testing (see [1], [2], [3], [5] and [9]). Ribenboim [10] and Granville [5], besides proving new results, provide a review of known facts and open problems.

By a classical Glaisher's result (see [4] or [7]) for a prime $p \geq 3$,

$$q_p(2) \equiv -\frac{1}{2}\sum_{k=1}^{p-1} \frac{2^k}{k} \pmod{p}. \tag{1.1}$$

Recently Skula conjectured that for any prime $p \geq 5$,

$$q_p(2)^2 \equiv -\sum_{k=1}^{p-1} \frac{2^k}{k^2} \pmod{p}. \tag{1.2}$$

2010 *Mathematics Subject Classification*: primary 11A07; secondary 11B65, 05A10.
Key words and phrases: congruence, Fermat quotient, harmonic numbers.
Received July 13, 2011, revised December 2011. Editor R. Kučera.
DOI: 10.5817/AM2012-2-113

Applying certain polynomial congruences, Granville [7] proved the congruence (1.2). In this note, we give an elementary proof of this congruence which is based on congruences for some harmonic type sums.

Remark 1.1. Recently, given a prime p and a positive integer $r < p-1$, R. Tauraso [14, Theorem 2.3] established the congruence $\sum_{k=1}^{p-1} 2^k/k^r \pmod p$ in terms of an alternating r-tiple harmonic sum. For example, combining this result when $r = 2$ with the congruence (1.2) [14, Corollary 2.4], it follows that

$$\sum_{1\le i<j\le p-1} \frac{(-1)^j}{ij} \equiv q_p(2)^2 \equiv -\sum_{k=1}^{p-1} \frac{2^k}{k^2} \pmod p .$$

2. Proof of the congruence (1.2)

The *harmonic numbers* H_n are defined by

$$H_n := \sum_{j=1}^{n} \frac{1}{j}, \qquad n = 1, 2, \dots ,$$

where by convention $H_0 = 0$.

Lemma 2.1. *For any prime $p \ge 5$ we have*

$$q_p(2)^2 \equiv \sum_{k=1}^{p-1} \Big(2^k + \frac{1}{2^k}\Big) \frac{H_k}{k+1} \pmod p . \tag{2.1}$$

Proof. In the present proof we will always suppose that i and j are positive integers such that $i \le p-1$ and $j \le p-1$, and that all the summations including i and j range over the set of such pairs (i, j).

Using the congruence (1.1) and the fact that by Fermat Little Theorem, $2^{p-1} \equiv 1 \pmod p$, we get

$$\begin{aligned} q_p(2)^2 &= \Big(\frac{2^{p-1}-1}{p}\Big)^2 \equiv \frac{1}{4}\Big(\sum_{k=1}^{p-1} \frac{2^k}{k}\Big)^2 = \frac{1}{4}\Big(\sum_{k=1}^{p-1} \frac{2^{p-k}}{p-k}\Big)^2 \\ &\equiv \frac{1}{4}\Big(2\sum_{k=1}^{p-1} \frac{2^{(p-1)-k}}{-k}\Big)^2 \equiv \Big(\sum_{k=1}^{p-1} \frac{1}{k\cdot 2^k}\Big)^2 \\ &= \sum_{i+j\le p} \frac{1}{ij\cdot 2^{i+j}} + \sum_{i+j\ge p} \frac{1}{ij\cdot 2^{i+j}} - \sum_{i+j=p} \frac{1}{ij\cdot 2^{i+j}} \pmod p . \end{aligned} \tag{2.2}$$

The last three sums will be called S_1, S_2 and S_3, respectively. We will determine them modulo p as follows.

$$\begin{aligned} S_1 &= \sum_{i+j\le p} \frac{1}{ij\cdot 2^{i+j}} = \sum_{k=2}^{p} \sum_{i+j=k} \frac{1}{ij\cdot 2^k} \\ &= \sum_{k=2}^{p} \frac{1}{2^k}\cdot\frac{1}{k}\sum_{i=1}^{k-1}\Big(\frac{1}{i}+\frac{1}{k-i}\Big) = \sum_{k=2}^{p} \frac{2H_{k-1}}{k\cdot 2^k} = \sum_{k=1}^{p-1} \frac{H_k}{(k+1)2^k} . \end{aligned} \tag{2.3}$$

Observe that the pair (i,j) satisfies $i+j=k$ for some $k\in\{p,p+1,\dots,2p-2\}$ if and only if for such a k holds $(p-i)+(p-j)=l$ with $l:=2p-k\le p$. Accordingly, using the fact that by Fermat Little Theorem, $2^{2p}\equiv 2^2 \pmod p$, we have

$$\begin{aligned}
S_2 &= \sum_{i+j\ge p}\frac{1}{ij\cdot 2^{i+j}} = \sum_{(p-i)+(p-j)\ge p}\frac{1}{(p-i)(p-j)\cdot 2^{(p-i)+(p-j)}}\\
&\equiv \sum_{i+j\le p}\frac{1}{ij\cdot 2^{2p-(i+j)}} \equiv \frac14\sum_{i+j\le p}\frac{2^{i+j}}{ij} = \frac14\sum_{k=2}^{p}\ \sum_{i+j=k}\frac{2^k}{ij}\\
&= \frac14\sum_{k=2}^{p}\frac{2^k}{k}\sum_{i=1}^{k-1}\Big(\frac1i+\frac{1}{k-i}\Big) = \sum_{k=2}^{p}\frac{2^{k-1}H_{k-1}}{k}\\
&= \sum_{k=1}^{p-1}\frac{2^kH_k}{k+1} \pmod p\,. \qquad (2.4)
\end{aligned}$$

By *Wolstenholme's theorem* (see, e.g., [15], [6]; for its generalizations see [11, Theorems 1 and 2]) if p is a prime greater than 3, then the numerator of the fraction $H_{p-1}=1+\frac12+\frac13+\cdots+\frac{1}{p-1}$ is divisible by p^2. Hence, we find that

$$\begin{aligned}
S_3 &= \sum_{i+j=p}\frac{1}{2^{i+j}ij} = \frac{1}{2^p}\sum_{i=1}^{p-1}\frac{1}{i(p-i)}\\
&= \frac{1}{p\cdot 2^p}\sum_{i=1}^{p-1}\Big(\frac1i+\frac{1}{p-i}\Big) = \frac{1}{p\cdot 2^{p-1}}H_{p-1}\equiv 0 \pmod p\,. \qquad (2.5)
\end{aligned}$$

Finally, substituting (2.3), (2.4) and (2.5) into (2.2), we immediately obtain (2.1). □

Proof of the following result easily follows from the congruence $H_{p-1}\equiv 0 \pmod p$.

Lemma 2.2 ([13, Lemma 2.1])**.** *Let p be an odd prime. Then*

$$H_{p-k-1}\equiv H_k \pmod p \qquad (2.6)$$

for every $k=1,2,\dots,p-2$.

Lemma 2.3. *For any prime $p\ge 5$ we have*

$$q_p(2)^2 \equiv \sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} - \sum_{k=1}^{p-1}\frac{2^k}{k^2} \pmod p\,. \qquad (2.7)$$

Proof. Since by Wolstenholme's theorem, $H_{p-1}/p \equiv 0 \pmod{p}$, using this and the congruences $2^{p-1} \equiv 1 \pmod{p}$ and (2.6) of Lemma 2.2, we immediately obtain

$$\begin{aligned}\sum_{k=1}^{p-1}\frac{2^kH_k}{k+1} &\equiv \sum_{k=1}^{p-2}\frac{2^kH_k}{k+1} = \sum_{k=1}^{p-2}\frac{2^{p-k-1}H_{p-k-1}}{p-k}\\ &\equiv -\sum_{k=1}^{p-2}\frac{H_k}{k\cdot 2^k} \equiv -\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} \pmod{p}. \qquad (2.8)\end{aligned}$$

Further, using Wolstenholme's theorem, we have

$$\begin{aligned}\sum_{k=1}^{p-1}\frac{H_k}{(k+1)2^k} &= 2\sum_{k=0}^{p-2}\frac{H_{k+1}-\frac{1}{k+1}}{(k+1)2^{k+1}} + \frac{H_{p-1}}{p\cdot 2^{p-1}}\\ &= 2\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} - 2\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k} + \frac{H_{p-1}}{p\cdot 2^{p-1}}\\ &\equiv 2\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} - 2\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k} \pmod{p}. \qquad (2.9)\end{aligned}$$

Moreover, from $2^p \equiv 2 \pmod{p}$ we have

$$\begin{aligned}\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k} &= \sum_{k=1}^{p-1}\frac{1}{(p-k)^2\cdot 2^{p-k}}\\ &\equiv \sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^{1-k}} = \frac{1}{2}\sum_{k=1}^{p-1}\frac{2^k}{k^2} \pmod{p}. \qquad (2.10)\end{aligned}$$

The congruences (2.8), (2.9) and (2.10) immediately yield

$$\begin{aligned}\sum_{k=1}^{p-1}\Big(2^k+\frac{1}{2^k}\Big)\frac{H_k}{k+1} &= \sum_{k=1}^{p-1}\frac{2^kH_k}{k+1} + \sum_{k=1}^{p-1}\frac{H_k}{(k+1)2^k}\\ &\equiv \sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} - \sum_{k=1}^{p-1}\frac{2^k}{k^2} \pmod{p}. \qquad (2.11)\end{aligned}$$

Finally, comparing (2.1) of Lemma 2.1 with (2.11), we obtain the desired congruence (2.7). □

Notice that the congruence $\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} \equiv 0 \pmod{p}$ with a prime $p \geq 5$ is recently established by Z.W. Sun [13, Theorem 1.1 (1.1)] and it is based on the identity from [13, Lemma 2.4]. Here we give another simple proof of this congruence (Lemma 2.6).

Lemma 2.4. *For any prime $p \geq 5$ we have*

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k} \equiv \frac{1}{2}\sum_{1\leq i\leq j\leq p-1}\frac{2^i-1}{ij} \pmod{p}. \qquad (2.12)$$

Proof. From the identity

$$\Big(\sum_{k=1}^{p-1}\frac{1}{k}\Big)\Big(\sum_{k=1}^{p-1}\frac{1}{k\cdot 2^k}\Big)=\sum_{1\le i<j\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{1\le j<i\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k},$$

and the congruence $H_{p-1}=1+\frac{1}{2}+\frac{1}{3}+\cdots+\frac{1}{p-1}\equiv 0\,(\mathrm{mod}\ p)$ it follows that

$$\sum_{1\le i<j\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{1\le j<i\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k}\equiv 0\pmod p. \tag{2.13}$$

Since $2^p\equiv 2\,(\mathrm{mod}\ p)$, we have

$$\sum_{1\le j<i\le p-1}\frac{1}{ij\cdot 2^j}\equiv\sum_{1\le j<i\le p-1}\frac{1}{2}\frac{2^{p-j}}{(p-i)(p-j)}\equiv\frac{1}{2}\sum_{1\le i<j\le p-1}\frac{2^j}{ij}\pmod p,$$

which substituting into (2.13) gives

$$\sum_{1\le i<j\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k}\equiv-\frac{1}{2}\sum_{1\le i<j\le p-1}\frac{2^j}{ij}\pmod p. \tag{2.14}$$

Further, if we observe that

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}=\sum_{k=1}^{p-1}\frac{H_{k-1}+\frac{1}{k}}{k\cdot 2^k}=\sum_{1\le i<j\le p-1}\frac{1}{ij\cdot 2^j}+\sum_{k=1}^{p-1}\frac{1}{k^2\cdot 2^k},$$

then substituting (2.14) into the previous identity, we obtain

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv-\frac{1}{2}\sum_{1\le i<j\le p-1}\frac{2^j}{ij}\pmod p. \tag{2.15}$$

Since

$$0\equiv\Big(\sum_{k=1}^{p-1}\frac{1}{k}\Big)\Big(\sum_{k=1}^{p-1}\frac{2^k}{k}\Big)=\sum_{1\le j\le i\le p-1}\frac{2^j}{ij}+\sum_{1\le i<j\le p-1}\frac{2^j}{ij}\pmod p,$$

comparing this with (2.15), we immediately obtain

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv\frac{1}{2}\sum_{1\le i\le j\le p-1}\frac{2^i}{ij}\pmod p. \tag{2.16}$$

From a well known fact that (see e.g., [9, p. 353])

$$\sum_{k=1}^{p-1}\frac{1}{k^2}\equiv 0\pmod p \tag{2.17}$$

we find that

$$\sum_{1\le i\le j\le p-1}\frac{1}{ij}=\frac{1}{2}\Big(\Big(\sum_{k=1}^{p-1}\frac{1}{k}\Big)^2+\sum_{k=1}^{p-1}\frac{1}{k^2}\Big)\equiv 0\pmod p.$$

Finally, the above congruence and (2.16) immediately yield the desired congruence (2.12). □

Lemma 2.5. *For any positive integer n we have*

$$\sum_{1\le i\le j\le n}\frac{2^i-1}{ij}=\sum_{k=1}^{n}\frac{1}{k^2}\binom{n}{k}. \tag{2.18}$$

Proof. Using the well known identities $\sum_{i=k}^{j}\binom{i-1}{k-1}=\binom{j}{k}$ and $\frac{1}{j}\binom{j}{k}=\frac{1}{k}\binom{j-1}{k-1}$ with $k\le j$, and the fact that $\binom{i}{k}=0$ when $i<k$, we have

$$\begin{aligned}
\sum_{1\le i\le j\le n}\frac{2^i-1}{ij}&=\sum_{1\le i\le j\le n}\frac{(1+1)^i-1}{ij}=\sum_{1\le i\le j\le n}\frac{1}{j}\sum_{k=1}^{i}\frac{1}{i}\binom{i}{k}\\
&=\sum_{1\le i\le j\le n}\frac{1}{j}\sum_{k=1}^{n}\frac{1}{k}\binom{i-1}{k-1}=\sum_{k=1}^{n}\frac{1}{k}\sum_{1\le i\le j\le n}\frac{1}{j}\binom{i-1}{k-1}\\
&=\sum_{k=1}^{n}\frac{1}{k}\sum_{k\le i\le j\le n}\frac{1}{j}\binom{i-1}{k-1}=\sum_{k=1}^{n}\frac{1}{k}\sum_{j=k}^{n}\frac{1}{j}\sum_{i=k}^{j}\binom{i-1}{k-1}\\
&=\sum_{k=1}^{n}\frac{1}{k}\sum_{j=k}^{n}\frac{1}{j}\binom{j}{k}=\sum_{k=1}^{n}\frac{1}{k}\sum_{j=k}^{n}\frac{1}{k}\binom{j-1}{k-1}\\
&=\sum_{k=1}^{n}\frac{1}{k^2}\sum_{j=k}^{n}\binom{j-1}{k-1}=\sum_{k=1}^{n}\frac{1}{k^2}\binom{n}{k},
\end{aligned}$$

as desired. □

Lemma 2.6 ([13, Theorem 1.1 (1.1)])**.** *For any prime $p\ge 5$ we have*

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv 0 \pmod p. \tag{2.19}$$

Proof. Using the congruence (2.12) from Lemma 2.4 and the identity (2.18) with $n=p-1$ in Lemma 2.5, we find that

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv\frac{1}{2}\sum_{k=1}^{p-1}\frac{1}{k^2}\binom{p-1}{k} \pmod p. \tag{2.20}$$

It is well known (see e.g., [8]) that for $k=1,2,\dots,p-1$,

$$\binom{p-1}{k}\equiv(-1)^k \pmod p. \tag{2.21}$$

Then from (2.20), (2.21) and (2.17) we get

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv\frac{1}{2}\sum_{k=1}^{p-1}\frac{(-1)^k}{k^2}=\frac{1}{2}\Big(2\sum_{\substack{1\le j\le p-1\\ 2|j}}\frac{1}{j^2}-\sum_{k=1}^{p-1}\frac{1}{k^2}\Big)$$

$$=\frac{1}{4}\sum_{k=1}^{(p-1)/2}\frac{1}{k^2}-\frac{1}{2}\sum_{k=1}^{p-1}\frac{1}{k^2}\equiv\frac{1}{4}\sum_{k=1}^{(p-1)/2}\frac{1}{k^2}\pmod p\,.$$

Finally, the above congruence together with the fact that from (2.17) (see e.g., [12, Corollary 5.2 (a) with $k=2$])

$$2\sum_{k=1}^{(p-1)/2}\frac{1}{k^2}\equiv\sum_{k=1}^{(p-1)/2}\frac{1}{k^2}+\sum_{k=1}^{(p-1)/2}\frac{1}{(p-k)^2}=\sum_{k=1}^{p-1}\frac{1}{k^2}\equiv 0\pmod p$$

yields

$$\sum_{k=1}^{p-1}\frac{H_k}{k\cdot 2^k}\equiv 0\pmod p\,.$$

This concludes the proof. □

Proof of the congruence (1.2)**.** The congruence (1.2) immediately follows from (2.7) of Lemma 2.3 and (2.19) of Lemma 2.6. □

References

[1] Agoh, T., Dilcher, K., Skula, L., *Fermat quotients for composite moduli*, J. Number Theory **66** (1997), 29–50.

[2] Cao, H. Q., Pan, H., *A congruence involving product of q–binomial coefficients*, J. Number Theory **121** (2006), 224–233.

[3] Ernvall, R., Metsänkylä, T., *On the p–divisibility of Fermat quotients*, Math. Comp. **66** (1997), 1353–1365.

[4] Glaisher, J. W. L., *On the residues of the sums of the inverse powers of numbers in arithmetical progression*, Quart. J. Math. **32** (1900), 271–288.

[5] Granville, A., *Some conjectures related to Fermat's Last Theorem*, Number Theory (Banff, AB, 1988), de Gruyter, Berlin (1990), 177–192.

[6] Granville, A., *Arithmetic properties of binomial coefficients. I. Binomial coefficients modulo prime powers*, Organic Mathematics–Burnaby, BC 1995, CMS Conf. Proc., vol. 20, Amer. Math. Soc., Providence, RI, 1997, pp. 253–276.

[7] Granville, A., *The square of the Fermat quotient*, Integers **4** (2004), # A22.

[8] Hardy, G. H., Wright, E. M., *An Introduction to the Theory of Numbers*, Clarendon Press, Oxford, 1960.

[9] Lehmer, E., *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Ann. Math. (1938), 350–360.

[10] Ribenboim, P., *13 lectures on Fermat's last theorem*, Springer–Verlag, New York, Heidelberg, Berlin, 1979.

[11] Slavutsky, I. Sh., *Leudesdorf's theorem and Bernoulli numbers*, Arch. Math. **35** (1999), 299–303.

[12] Sun, Z. H., *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (1–3) (2000), 193–223.

[13] Sun, Z. W., *Arithmetic theory of harmonic numbers*, Proc. Amer. Math. Soc. **140** (2012), 415–428.

[14] Tauraso, R., *Congruences involving alternating multiple harmonic sums*, Electron. J. Comb. **17** (2010), # R16.

[15] Wolstenholme, J., *On certain properties of prime numbers*, Quart. J. Pure Appl. Math. **5** (1862), 35–39.

Maritime Faculty, University of Montenegro,
Dobrota 36, 85330 Kotor, Montenegro
E-mail: romeo@ac.me