ARCHIVUM MATHEMATICUM (BRNO)
Tomus 53 (2017), 101–109

ON THE EXISTENCE OF NON-LINEAR FRAMES

Shah Jahan, Varinder Kumar, and S.K. Kaushik

Abstract. A stronger version of the notion of frame in Banach space called Strong Retro Banach frame (SRBF) is defined and studied. It has been proved that if $\mathcal{X}$ is a Banach space such that $\mathcal{X}^*$ has a SRBF, then $\mathcal{X}$ has a Bi-Banach frame with some geometric property. Also, it has been proved that if a Banach space $\mathcal{X}$ has an approximative Schauder frame, then $\mathcal{X}^*$ has a SRBF. Finally, the existence of a non-linear SRBF in the conjugate of a separable Banach space has been proved.

1. Introduction

Frames for Hilbert spaces were introduced by Duffin and Schaeffer [5] in the context of nonharmonic Fourier series. Frames now a days are widely used in various branches of mathematics and engineering. Feichtinger and Grochenig [6] generalized the notion of frame to Banach spaces and introduced the concept of atomic decomposition in a Banach space. Also, Grochenig [7] introduced a more general concept namely Banach frame for Banach spaces. For a nice and comprehensive survey of frames and related concepts one may refer to [1, 4].

Various other generalizations of frames for Banach spaces were defined and studied by many authors namely Schauder frames by Han and Larson [8] and also studied by Casazza et al. [2, 3], frames by Terekhin [17]. Banach frames in conjugate Banach spaces, called retro Banach frames, were introduced and studied by Jain et al. [9] and further studied in [13]. Approximative atomic decompositions in Banach spaces were studied in [10]. Schauder frames in conjugate Banach spaces were defined and studied in [12] while approximative Schauder frames were studied in [11]. The notion of Bi-Banach frame in a Banach space was defined and studied in [14] wherein they noted that a Schauder frame for a Banach space is a Bi-Banach frame but the converse is not true.

In the present paper, we shall consider a stronger notion of frame in a Banach space called strong Retro Banach frame (SRBF). It has been proved that if $\mathcal{X}$ is a Banach space such that $\mathcal{X}^*$ has a SRBF, then $\mathcal{X}$ has a Bi- Banach frame with some geometric property. Also, it has been proved that if a Banach space $\mathcal{X}$ has an approximative Schauder frame, then $\mathcal{X}^*$ has a SRBF. Finally, a result related to

2010 *Mathematics Subject Classification*: primary 42C15; secondary 46B15.
Key words and phrases: Banach frames, retro Banach frames, approximative Schauder frames.
The research work of the first author is partially supported by UGC, India.
Received November 2, 2015, revised February 2017. Editor W. Kubiś.
DOI: 10.5817/AM2017-2-101

the existence of a non-linear SRBF in the conjugate of a separable Banach space has been proved.

Throughout this paper $\mathcal{X}$ will denotes an infinite dimensional Banach space over the scalar field $\mathbb{K}(\mathbb{R},\mathbb{C})$, $\mathcal{X}^*$ denotes the conjugate space of $\mathcal{X}$ and $L(\mathcal{X},\mathcal{X})$ denote the Banach space of all continuous linear mappings of $\mathcal{X}$ into $\mathcal{X}$. For a sequence $\{x_n\}\in\mathcal{X}$ and $\{f_n\}\in\mathcal{X}^*$, $[x_n]$ denotes the closed linear span of $\{x_n\}$ in the norm topology of $\mathcal{X}$ and $\overline{[f_n]}$ the closed linear span of $\{f_n\}$ in the weak star topology of $\mathcal{X}^*$. A sequence space S is called a *BK-space* if it is a Banach space and the co-ordinate functionals are continuous on S. That is the relations $x_n=\{\alpha_j{}^{(n)}\}$, $x=\{\alpha_j\}\in S$, $\lim\limits_{n\to\infty}x_n=x$ imply $\lim\limits_{n\to\infty}\alpha_j^{(n)}=\alpha_j$ $(j=1,2,3,\dots)$. Also, if $V\subseteq\mathcal{X}^*$, then we define $\gamma(v)=\inf\limits_{\substack{x\in\mathcal{X}\\ x\neq0}}\sup\limits_{\substack{f\in v\\ \|f\|\leq I}}\left|f\left(\frac{x}{\|x\|}\right)\right|$.

A sequence $\{x_n\}\subset\mathcal{X}$ is said to be a *Markusevic basis* (M-basis) for $\mathcal{X}$ if $\{x_n\}$ is complete in $\mathcal{X}$ and there exists a sequence $\{f_n\}$ in $\mathcal{X}^*$ biorthogonal to $\{x_n\}$, called an associated sequence of coefficient functional (a.s.c.f.), which is total on $\mathcal{X}$.

Definition 1.1 ([9])**.** Let $\mathcal{X}$ be a Banach space and $\mathcal{X}_d$ be a BK-space. Let $\{x_n\}\subset\mathcal{X}$ and $J\colon\mathcal{X}_d^*\to\mathcal{X}^*$ be given. The pair $(\{x_n\},J)$ is called a *retro Banach frame* for $\mathcal{X}^*$ with respect to $\mathcal{X}_d^*$ if

(a) $\{f(x_n)\}\in\mathcal{X}_d^*$, for all $f\in\mathcal{X}^*$.

(b) There exist positive constants A and B with $0<A\leq B<\infty$ such that

$$A\|f\|_{\mathcal{X}^*}\leq\|\{f(x_n)\}\|_{\mathcal{X}_d^*}\leq B\|f\|_{\mathcal{X}^*},\quad\text{for all}\quad f\in\mathcal{X}^*. \tag{1.1}$$

(c) J is a bounded linear operator such that

$$J(\{f(x_n)\})=f,\quad\text{for all}\quad f\in\mathcal{X}^*.$$

The constants A and B are called lower and upper bounds of the retro Banach frame $(\{x_n\},J)$. The inequality (1.1) is called the retro Banach frame inequality.

A retro Banach frame $(\{x_n\},J)$ is said to be exact if there exists a sequence $\{f_n\}\subset\mathcal{X}^*$ such that $f_i(x_j)=\delta_{i,j}$, for all $i,j\in\mathbb{N}$.

Definition 1.2. Let $\mathcal{X}$ be a Banach space with dual $\mathcal{X}^*$. A pair $(\{x_n\},\{f_n\})$ (where $\{f_n\}\subset\mathcal{X}^*$ and $\{x_n\}\subset\mathcal{X}$) is called a *Bi-Banach frame* for $\mathcal{X}$ if there exist associated Banach spaces $\mathcal{X}_d$ and $(\mathcal{X}^*)_d$ and bounded linear operators $S\colon\mathcal{X}_d\to\mathcal{X}$, $T\colon(\mathcal{X}^*)_d\to\mathcal{X}^*$ such that $(\{f_n\},S)$ is a Banach frame for $\mathcal{X}$ and $(\{x_n\},T)$ is retro Banach frame for $\mathcal{X}^*$.

A Bi-Banach frame $(\{x_n\},\{f_n\})$ is called *tight* if both the retro Banach frame $(\{x_n\},T)$ and the Banach frame $(\{f_n\},S)$ are tight.

The following results are stated in the form of lemmas which will be used in the subsequent work.

Lemma 1.3 ([16])**.** *Let $\mathcal{X}$ be a Banach space and $\{f_n\}\subset\mathcal{X}^*$ be a sequence such that $\{x\in\mathcal{X}:f_n(x)=0,\forall\ n\in\mathbb{N}\}=\{0\}$. Then $\mathcal{X}$ is linearly isometric to the Banach space $\mathcal{X}_d=\{\{f_n(x)\}:x\in\mathcal{X}\}$, where the norm is given by $\|\{f_n(x)\}\|_{\mathcal{X}_d}=\|x\|_{\mathcal{X}},x\in\mathcal{X}$.*

Lemma 1.4 ([16]). *Let $\mathcal{X}$ be a separable normed linear space and let $\{x_n^*\}$ be a sequence in $\mathcal{X}^*$ such that $\frac{x_n^*}{\|x_n^*\|} \xrightarrow{w^*} 0$ and that for the linear subspace $[x_n^*]$ of $\mathcal{X}^*$, $\gamma([x_n^*]) \geq 0$. Then there exist a norm $|\cdot|$ on $\mathcal{X}$ equivalent to the initial norm on $\mathcal{X}$ such that $(\mathcal{X}, |\cdot|)$ is strictly convex and satisfies the following property*

$$(1.2) \qquad \text{If} \quad \lim_{n\to\infty} f_k(x_n) = f_k(x_0)(k = 1, 2, \dots), \quad \text{then} \quad \lim_{n\to\infty} |x_n| \geq |x_0|.$$

2. Main result

Approximative Schauder frames in Banach spaces were studied in [11] and the notion of Bi-Banach frame was studied in [14]. In the following definition, we gave a stronger notion called Strong Retro Banach frame (SRBF). The idea of defining this notion is to correlate this notion with the existing notions like approximative Schauder frames and Bi-Banach frames.

Definition 2.1. Let $\{x_n\} \subset \mathcal{X}$ be an exact RBF for $\mathcal{X}^*$ with admissible sequence $\{f_n\} \subset \mathcal{X}^*$. Let $X_n = [x_1, x_2, \dots, x_n]$, $n \in \mathbb{N}$. If there exists a sequence $\{v_n\}$, where each $v_n \colon X_n \to X_n$ is a continuous linear mapping, such that $x = \lim_{n\to\infty} v_n \sum_{i=1}^{n} f_i(x)x_i$, $x \in \mathcal{X}$, then $(\{x_n\}, \{f_n\}, \{v_n\})$ is called a strong RBF (or SRBF) for $\mathcal{X}^*$.

Remark 2.2. If we define $u_n \colon \mathcal{X} \to \mathcal{X}$, $n \in \mathbb{N}$ by

$$u_n(x) = v_n \sum_{i=1}^{n} f_i(x)x_i, \quad n \in \mathbb{N}.$$

Then one may observe that if $(\{x_n\}, \{f_n\}, \{v_n\})$ is a SRBF, then $\lim_{n\to\infty} u_n(x) = x$ and $\dim u_n(\mathcal{X}) = \dim (v_n \sum_{i=1}^{n} f_i(x)x_i)(\mathcal{X}) \leq n < \infty$ and so $\{x_n\}$ is an approximative basis of $\mathcal{X}$.

In the following result, we prove that the existence of SRBF in the conjugate of a Banach space guarantees the existence of a Bi-Banach frame in the Banach space along with some geometric property.

Theorem 2.3. *Let X be a Banach space and $(\{x_n\}, \{f_n\}, \{v_n\})$ be a SRBF for $\mathcal{X}^*$ with admissible sequence $\{f_n\} \subset \mathcal{X}^*$. Then $(\{x_n\}, \{f_n\})$ is a Bi-Banach frame for $\mathcal{X}$ such that $\gamma([f_n]) > 0$.*

Proof. Clearly, by definition of SRBF, $\{x \in \mathcal{X} : f_n(x) = 0, \text{for all } n \in \mathbb{N}\} = \{0\}$. Therefore, by Lemma 1.3, there exists an associated Banach space $\mathcal{X}_d = \{\{f_n(x)\}; x \in \mathcal{X}\}$ with norm given by $\|\{f_n(x)\}\|_{\mathcal{X}_d} = \|x\|_{\mathcal{X}}$, $x \in \mathcal{X}$. Define $J \colon \mathcal{X}_d \to \mathcal{X}$ by $J(\{f_n(x)\}) = x$, $x \in \mathcal{X}$. Then J is a bounded linear operator such that $(\{f_n\}, J)$ is a Banach frame for $\mathcal{X}$. Hence $(\{x_n\}, \{f_n\})$ is a Bi-Banach frame for $\mathcal{X}$. Let for each $n \in \mathbb{N}$, $v_n \colon X_n \to X_n$ be a continuous linear mapping given by $\lim_{n\to\infty} v_n \sum_{i=1}^{n} f_i(x)x_i = x$. Write $v_n(x_j) = \sum_{i=1}^{n} a_{ji}^{(n)} x_i$, $j = 1, 2, \dots, n$, $n \in \mathbb{N}$, where

$a_{ji}^{(n)} = f_i(v_n(x_j))$, for all $i, j = 1, 2, 3, \ldots, n$, $n \in \mathbb{N}$. Thus

$$v_n\Big(\sum_{i=1}^{n} f_i(x)x_i\Big) = \sum_{j=1}^{n}\Big(\sum_{i=1}^{n} a_{ij}^{(n)} f_i(x)\Big)x_j\,, \quad x \in \mathcal{X},\ n \in \mathbb{N}\,.$$

Define

$$h_{n,j} = \sum_{i=1}^{n} a_{ij}^{(n)} f_i\,, \quad j = 1, 2, 3, \ldots, n,\ n \in \mathbb{N}\,.$$

Then $h_{n,i} \in [f_i]_{i=1}^{n}$ $(j = 1, 2, \ldots, n;\ n \in \mathbb{N})$. Hence, we conclude that $\gamma([f_n]) > 0$. □

In the following result, we prove a weak duality type result.

Theorem 2.4. *Let $(\{x_n\}, \{f_n\}, \{v_n\})$ be a SRBF for $\mathcal{X}^*$ with admissible sequence $\{f_n\} \subset \mathcal{X}^*$. Then there exists a sequence of continuous linear mappings $\{\tau_n\}$ ($\tau_n \colon V_n \to V_n$, where $V_n = [f_1, f_2, \ldots, f_n], n \in \mathbb{N}$) such that*

$$f(x) = \lim_{n\to\infty}\Big(\tau_n \sum_{i=1}^{n} f(x_i)f_i\Big)(x)$$

Proof. For each $k = 1, 2, 3, \ldots, n$, $n \in \mathbb{N}$, define $\tau_n \colon \operatorname{span}\{f_1, f_2, \ldots, f_n\} \to \operatorname{span}\{f_1, f_2, \ldots, f_n\}$ by

$$\tau_n(f_k) = \sum_{i=1}^{n} a_{ik}^{(n)} f_i = \sum_{i=1}^{n} f_k(v_n(x_i))f_i\,.$$

Extend each τ_n to $[f_1, f_2, \ldots, f_n]$. Then

$$\begin{aligned}
\Big(\tau_n \sum_{i=1}^{n} f(x_i)f_i\Big)(x) &= \sum_{i=1}^{n} f(x_i)\big(\tau_n(f_i)\big)(x) \\
&= \sum_{i=1}^{n}\Big(\sum_{j=1}^{n} a_{ji}^{(n)} f_j(x)\Big)f(x_i) \\
&= \sum_{i=1}^{n}\Big(\sum_{j=1}^{n} f_i\big(v_n(x_j)\big)f_j(x)\Big)f(x_i) \\
&= \sum_{i=1}^{n} v_n\Big(\sum_{j=1}^{n} f_i(x_j)f_j(x)\Big)f(x_i) \\
&= f\Big(v_n\Big(\sum_{i=1}^{n} f_i(x)x_i\Big)\Big) \\
&= \to f(x) \quad \text{as} \quad n \to \infty
\end{aligned}$$

□

Approximative Schauder frames were defined and studied in [11]. In the following result, we prove that if a Banach space $\mathcal{X}$ has an approximative Schauder frame, then its dual space has a SRBF.

Theorem 2.5. *If a Banach space $\mathcal{X}$ has an approximative Schauder frame, then $\mathcal{X}^*$ has a SRBF.*

Proof. Let $\{u_n\}$ be a sequence of finite rank continuous linear mapping from $\mathcal{X}$ to $\mathcal{X}$ such that $\lim_{n\to\infty} u_n(x) = x$, $x \in \mathcal{X}$. Let $\{x_n\}$ be a Markusevic basis for $\mathcal{X}$ with a.s.c.f. $\{f_n\} \subset \mathcal{X}^*$ such that

$$\bigcup_n u_n^*(\mathcal{X}^*) \subset [f_n]\,. \tag{2.1}$$

Since each u_n is finite dimensional, we may write

$$u_n(x) = \sum_{i=1}^{p_n} \psi_{ni}(x)\phi_{ni}\,, \qquad x \in \mathcal{X},\ n \in \mathbb{N}\,,$$

where $\{\phi_{ni}\}_{i=1}^{p_n}$ is a basis for $\{u_n(\mathcal{X})\}$ with associated sequence $\{\psi_{ni}\}_{i=1}^{p_n} \subset \mathcal{X}^*$. Let $\{g_{ni}\}_{i=1}^{p_n}$ be a sequence in $\mathcal{X}^*$ that is biorthogonal to $\{\phi_{ni}\}_{i=1}^{p_n}$. Then

$$\begin{aligned} u_n^*(g_{nj})(x) &= g_{nj}\big(u_n(x)\big) \\ &= g_{nj}\Big(\sum_{i=1}^{p_n} \psi_{ni}(x)\phi_{n_i}\Big) \\ &= \sum_{i=1}^{p_n} \psi_{ni}(x) g_{nj}(\phi_{n_i}) \\ &= \psi_{nj}(x)\,, \qquad j = 1, 2, \ldots, p_n\,. \end{aligned}$$

Hence, $\psi_{nj} \in [f_n]$, $j = 1, 2, \ldots, p_n$, $n \in \mathbb{N}$. Let $n \in \mathbb{N}$ be given. Then, for any $\epsilon > 0$, there exists an integer $m_n(\epsilon)$ such that for each $i = 1, 2, \ldots, p_n$ one can find $\overline{\phi}_{ni} \in [x_1, x_2, \ldots, x_{m_n}]$ and $\overline{\psi}_{ni} \in [f_1, f_2, \ldots, f_{m_n}]$ such that

$$\|\phi_{ni} - \overline{\phi}_{ni}\| < \epsilon \quad \text{and} \quad \|\psi_{ni} - \overline{\psi}_{ni}\| < \epsilon\,, \qquad i = 1, 2, 3, \ldots, p_n\,. \tag{2.2}$$

Write

$$\overline{v}_{m_n}(x) = \sum_{i=1}^{p_n} \overline{\psi}_{ni}(x)\overline{\phi}_{ni}\,, \qquad x \in \mathcal{X}\,.$$

Then

$$\begin{aligned} \|\overline{v}_{m_n}(x) - u_n(x)\| &= \Big\| \sum_{i=1}^{p_n} \big(\overline{\psi}_{ni}(x) - \psi_{ni}(x)\big)\overline{\phi}_{ni} + \sum_{i=1}^{p_n} \psi_{ni}(x)(\overline{\phi}_{ni} - \phi_{ni})\Big\| \\ &\le \Big(\sum_{i=1}^{p_n} \|\overline{\psi}_{ni} - \psi_{ni}\|\|\overline{\phi}_{ni}\| + \sum_{i=1}^{p_n} \|\psi_{ni}\|\|\overline{\phi}_{ni} - \phi_{ni}\|\Big) \|x\|\,,\ x \in \mathcal{X}\,. \end{aligned}$$

Therefore, by (2.2), taking $\epsilon = \frac{1}{n}$ and $\{m_n\}$ to be an increasing sequence, we obtain

$$\|\overline{v}_{m_n} - u_n\| < \frac{1}{n}\,. \tag{2.3}$$

Now, observe that

$$f_i\Big(x-\sum_{j=1}^{k} f_j(x)x_j\Big)=0\,, \tag{2.4}$$
$$\text{for all}\quad x\in\mathcal{X},\ i=1,2,\dots,m_n\,;\quad k=m_n,m_n+1,\dots.$$

Also, $\overline{\psi}_{ni}\in[f_1,f_2,\dots,f_{m_n}]$. So, for $x\in\mathcal{X}$, we have

$$\begin{aligned}\overline{v}_{m_n}\Big(\sum_{j=1}^{k} f_j(x)x_j\Big)&=\sum_{i=1}^{p_n}\overline{\psi}_{ni}\Big(\sum_{j=1}^{k} f_j(x)x_j\Big)\overline{\phi}_{ni}\\&=\sum_{i=1}^{p_n}\overline{\psi}_{ni}(x)\overline{\phi}_{ni}\\&=\overline{v}_{m_n}(x)\,,\quad\text{for all}\quad x\in\mathcal{X}\quad\text{and}\quad k\geq m_n\,.\end{aligned}\tag{2.5}$$

Therefore

$$\begin{aligned}\lim_{n\to\infty}\overline{v}_{m_n}\Big(\sum_{j=1}^{m_n} f_j(x)x_j\Big)&=\lim_{n\to\infty}\overline{v}_{m_n}(x)\\&=\lim_{n\to\infty}u_n(x)\\&=x\,,\qquad x\in\mathcal{X}\,.\end{aligned}\tag{2.6}$$

Define a sequence $\{T_n\}$ by $T_n(x)=\sum_{i=1}^{n} f_i(x)x_i,\ \ n\in\mathbb{N}$. Write

$$v_k=T_k|_{[x_1,x_2,\dots,x_k]}\,,\quad k=1,2,\dots,m_1-1$$

and

$$v_k=\overline{v}_{m_n}|_{[x_1,x_2,\dots,x_k]}\,,\quad k=m_n,m_n+1,\dots,m_n-1\,,\qquad n=1,2,3\dots\ .$$

Then, each v_k is a continuous linear mapping defined on $[x_1,x_2,\dots,x_k]$ with range given by

$$v_k\big([x_1,x_2,\dots,x_k]\big)=[x_1,x_2,\dots,x_k]\,;\quad k=1,2,3,\dots,m_1-1$$

$$v_k\big([x_1,x_2,...x_k]\big)\subset[x_1,x_2,\dots,x_k]\,,\quad k=m_n,m_n+1,\dots,m_{n+1}-1\,,\quad n\in\mathbb{N}\,.$$

Using (2.5) and (2.6), we obtain $\lim_{n\to\infty}\overline{v}_{m_n}(x)=x$. Hence $\lim_{n\to\infty}v_n\big(\sum_{i=1}^{n} f_i(x)x_i\big)=x$. □

In view of the proof of Theorem 2.5, one may observe that existence of approximative Schauder frame in a Hilbert space is a sufficient condition for the space having Markusevic basis to have a SRBF. More precisely we have

Corollary 2.6. *Let $\mathcal{H}$ be a Hilbert space with an approximative Schauder frame. Then every Markusevic basis of $\mathcal{H}$ give rise to a SRBF for $\mathcal{H}$.*

In the following example, we show that in general, a SRBF do not have strong duality

Example 2.7. Let $\mathcal{X}$ be a Banach space with a Schauder basis and such that $\mathcal{X}^*$ is separable but fails to have approximative property. Let $\{x_n\}$ be a shrinking Markusevic basis of $\mathcal{X}$ with associated sequence of coefficient functional $\{f_n\} \subset \mathcal{X}^*$. Define

$$u_n(x) = \sum_{i=1}^{n} f_i(x)x_i\,, \qquad x \in \mathbb{N}\,.$$

Then $\{x_n\}$ is an approximative Schauder frame for $\mathcal{X}$ satisfying $\bigcup_{n=1}^{\infty} {u_n}^*(\mathcal{X}^*) \subset [f_n]$. Therefore $(\{x_n\}, \{f_n\}, \{v_n\})$ is a SRBF for $\mathcal{X}^*$. However, $\mathcal{X}^{**}$ has no SRBF.

One may observe that in Definition 2.1, each v_n is linear. Now, we would like to drop this condition of linearity and in the process define non-linear SRBF.

Definition 2.8. A SRBF $(\{x_n\}, \{f_n\}, \{v_n\})$ is called non-linear SRBF if each v_n is continuous but not necessarily linear.

Finally, we prove the following result related to the existence of a non-linear SRBF.

Theorem 2.9. *If $\mathcal{X}$ is a separable Banach Space, then $\mathcal{X}^*$ has a non-linear SRBF.*

Proof. Let $\{x_n\}$ be a Markusevic basis with a sequence of coefficient functional $\{f_n\} \subset \mathcal{X}^*$ such that $\gamma([f_n]) > 0$. Then, by Lemma 1.4, there is a norm $|\cdot|$ on $\mathcal{X}$ that is equivalent to the original norm $\|\cdot\|$ on $\mathcal{X}$ such that $\mathcal{X}$ with this new norm $|\cdot|$ is strictly convex. Therefore, by [15, Corollary 3.3, page 110], for every finite dimensional subspace $\mathcal{G}$ of $\mathcal{X}$ and for every $x \in \mathcal{X} \setminus \mathcal{G}$, there is a unique $\pi_{\mathcal{G}}(x) \in \mathcal{G}$ such that $|x - \pi_{\mathcal{G}}(x)| = \operatorname{dist}(x, \mathcal{G}) = \min\limits_{x \in \mathcal{G}} |x - g|$ and such that the mapping $\pi_{\mathcal{G}} \colon \mathcal{X} \to \mathcal{G}$ is continuous (here note that, in general, $\pi_{\mathcal{G}}$ is non-linear). Let $\mathfrak{N}(a, b)$ denote a positive integer depending on a and b. For each n, choose an increasing sequence of positive integers $\{m_n\}$ with $m_1 = \mathfrak{N}(1, 1)$, $m_2 = \mathfrak{N}(m_1, \frac{1}{2})$, $m_3 = \mathfrak{N}(m_2, \frac{1}{3})$, $\ldots$, $m_n = \mathfrak{N}(m_{n-1}, \frac{1}{n})$, for all $n \geq 2$ and satisfying

$$\operatorname{dist}\,\big(a, [x_i]_{i=m_{n-1}+1}^{m_n}\big) \leq \Big(1 + \frac{1}{n}\Big) \operatorname{dist}\,\big(a, [x_i]_{i=m_{n-1}+1}^{\infty}\big)\,,$$

where $a \in [x_i]_{i=1}^{m_{n-1}}$. Define $\{v_n\}$ by $v_k = T_k|_{[x_1, \ldots, x_k]}$, $k = 1, 2, \ldots, m_1 - 1$, where $T_k(x) = \sum_{i=1}^{k} f_i(x)x_i$ and for any $b = \sum_{i=1}^{k} a_i x_i \in [x_i]_{i=1}^{k}$, $(k = m_n, m_n + 1, \ldots, m_{n+1} - 1;\ n \in \mathbb{N})$

$$v_k(b) = \sum_{i=1}^{m_{n-1}} a_i x_i - \pi_{\mathcal{G}}\Big(\sum_{i=1}^{m_{n-1}} a_i x_i\Big)\,,$$

where $\mathcal{G} = [x_i]_{i=m_{n-1}+1}^{m_n}$. Then each v_n is continuous (in general, non-linear) with range given by

$$v_k\big([x_1, \ldots, x_k]\big) = [x_1, \ldots, x_k]\,, \quad k = 1, 2, 3, \ldots, m_1 - 1$$

$$v_k\big([x_1, \ldots, x_k]\big) \subset [x_1, \ldots, x_k]\,, \quad (k = m_n, m_n + 1, \ldots, m_{n+1} - 1;\ n \in \mathbb{N})\,.$$

Let $x \in \mathcal{X}$ be any element. Then

$$f_i\Big(v_k\Big(\sum_{j=1}^{k} f_j(x)x_j\Big)\Big) = f_i(x)\,,\ i = 1, 2, \ldots, m_n\,;\ k = m_n, m_n{+}1, \ldots, m_{n+1}{-}1; n \in \mathbb{N}\,.$$

This gives

$$\lim_{k\to\infty} f_i\Big(v_k\Big(\sum_{j=1}^{k} f_j(x)x_j\Big)\Big) = f_i(x)\,, \quad i = 1, 2, \ldots\,. \tag{2.7}$$

In view of Lemma 1.4, we have

$$\lim_{k\to\infty} \Big|v_k\Big(\sum_{i=1}^{k} f_i(x)x_i\Big)\Big| \geq |x|\,. \tag{2.8}$$

Also, we have

$$\begin{aligned}
v_k\Big(\sum_{i=1}^{k} f_i(x)x_i\Big) &= \Big|\sum_{i=1}^{m_{n-1}} f_i(x)x_i - \pi_{\mathcal{G}} \sum_{i=1}^{m_{n-1}} f_i(x)x_i\Big| \\
&= \operatorname{dist}\Big(\sum_{i=1}^{m_{n-1}} f_i(x)x_i, \mathcal{G}\Big) \\
&\leq \Big(1 + \frac{1}{n}\Big) \operatorname{dist}\Big(\sum_{i=1}^{m_{n-1}} f_i(x)x_i, [x_i]_{i=m_{n-1}+1}^{\infty}\Big) \\
&\leq \Big(1 + \frac{1}{n}\Big)\Big|\sum_{i=1}^{m_{n-1}} f_i(x)x_i + \Big(x - \sum_{i=1}^{m_{n-1}} f_i(x)x_i\Big)\Big| \\
&= \Big(1 + \frac{1}{n}\Big)|x|\,, \quad k = m_n, m_n + 1, \quad m_{n+1} - 1;\ n \in \mathbb{N}\,.
\end{aligned}$$

Thus, by (2.8), we have

$$\lim_{k\to\infty} \Big|v_k\Big(\sum_{i=1}^{k} f_i(x)x_i\Big)\Big| = |x|\,. \tag{2.9}$$

Hence, we conclude that

$$\lim_{k\to\infty} \Big|v_k\Big(\sum_{i=1}^{k} f_i(x)x_i\Big) - x\Big| = 0\,.$$

Since $|\cdot|$ is equivalent to the initial norm of $\mathcal{X}$, we obtain

$$\lim_{n\to\infty} v_n\Big(\sum_{i=1}^{n} f_i(x)x_i\Big) = x\,, \quad x \in \mathcal{X}\,.$$

□

Acknowledgement. The authors would like to give thanks to the referee for his critical remarks and suggestions for the improvement of the paper.

References

[1] Casazza, P.G., *The art of frame theory*, Taiwanese J. Math. **4** (2000), 129–201.

[2] Casazza, P.G., Dilworth, S.J., Odell, E., Th.Schlumprecht, Zsak, A., *Cofficient quantization for frames in Banach spaces*, J. Math. Anal. Appl. **348** (2008), 66–86.

[3] Casazza, P.G., Han, D., Larson, D., *Frames for Banach spaces*, Contemp. Math. **247** (1999), 149–181.

[4] Christensen, O., *Frames and bases (An introductory course)*, Birkhäuser, Boston, 2008.

[5] Duffin, R., Schaeffer, A., *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc. **72** (1952), 341–366.

[6] Feichtinger, H.G., Grochenig, K.H., *A unified approach to atomic decompositions via integrable group representations*, Lecture Notes in Math., vol. 1302, Springer, 1988, pp. 52–73.

[7] Grochenig, K.H., *Describing functions: Atomic decompositions versus frames*, Monatsh. Math. **112** (1991), 1–41.

[8] Han, D., Larson, D.R., *Frames, bases and group representations*, Mem. Amer. Math. Soc. **147** (2000), 1–91.

[9] Jain, P.K., Kaushik, S.K., Vashisht, L.K., *Banach frames for conjugate Banach spaces*, Z. Anal. Anwendungen **23** (2004), 713–720.

[10] Kaushik, S.K., Sharma, S.K., *On approximative atomic decompositions in Banach spaces*, Communications in Mathematics and Applications **3** (3) (2012), 293–301.

[11] Kaushik, S.K., Sharma, S.K., *Generalized Schauder frames*, Arch. Math. (Brno) **50** (2014), 39–49.

[12] Kaushik, S.K., Sharma, S.K., Poumai, K.T., *On Schauder frames in conjugate Banach spaces*, Journal of Mathematics **2013** (2013), 4 pages, Article ID 318659.

[13] Poumai, K.T., Kaushik, S.K., *Retro Banach frames, almost exact retro Banach frames in Banach spaces*, Bulletin Math. Anal. Appl. **7** (1) (2015), 38–48.

[14] Sharma, S.K., *On Bi-Banach frames in Banach spaces*, Int. J. Wavelets Multiresolut Inf. Process **12** (2) (2014), 10 pages, 1450015.

[15] Singer, I., *Best approximation in normed linear spaces by elements of linear subspaces*, Springer-Verlag , New York, Heidelberg, Berlin, 1970.

[16] Singer, I., *Bases in Banach spaces II*, Springer-Verlag, New York, Heidelberg, Berlin, 1981.

[17] Terekhin, P.A., *Frames in Banach spaces*, Funct. Anal. Appl. **44** (3) (2010), 199–208.

S. Jahan, Department of Mathematics,
University of Delhi, Delhi–110 007, India
E-mail: chowdharyshahjahan@gmail.com

V. Kumar, Department of Mathematics,
Shaheed Bhagat Singh College,
University of Delhi, Delhi–110 017, India
E-mail: varinder1729@gmail.com

S.K. Kaushik, Department of Mathematics,
Kirori Mal College,
University of Delhi, Delhi–110 007, India
E-mail: shikk2003@yahoo.co.in