ARCHIVUM MATHEMATICUM (BRNO)
Tomus 54 (2018), 111–117

SOME FUNCTORIAL PROLONGATIONS OF GENERAL CONNECTIONS

Ivan Kolář

Abstract. We consider the problem of prolongating general connections on arbitrary fibered manifolds with respect to a product preserving bundle functor. Our main tools are the theory of Weil algebras and the Frölicher-Nijenhuis bracket.

0. Introduction

Our approach to connections on an arbitrary fibered manifold $p\colon Y \to M$ is slightly different from the approach by C. Ehresmann, [2], p. 186. Roughly speaking, the fundamental idea in [2] is the development along the individual curves, while the main idea of our approach is the absolute differentiation of the sections of Y. This is explained in Chapter 1 of the present paper. But the theory of general connections on Y can be well developed even by using the concept of tangent valued form on Y. This was invented by L. Mangiarotti and M. Modugno in [7] and first systematically presented in the book [6]. We repeat the basic ideas in Chapter 2. Chapter 3 is devoted to the case of product preserving bundle functors on the category $\mathcal{M}f$ of smooth manifolds and smooth maps. Our geometrical description of them uses the language of Weil algebras, [5], [6].

All manifolds and maps are assumed to be infinitely differentiable. Unless otherwise specified, we use the terminology and notation from [6].

1. General connections

Let $\pi_Y\colon TY \to Y$ denote the tangent bundle of a fibered manifold $p\colon Y \to M$. In [6], a general connection of Y is defined as a lifting map

$$\Gamma\colon Y \times_M TM \to TY \tag{1}$$

linear in TM and satisfying $\pi_Y \circ \Gamma = pr_1$, $Tp \circ \Gamma = pr_2$, $Y \xleftarrow{pr_1} Y \times_M TM \xrightarrow{pr_2} TM$. If x^i, y^p are some local fiber coordinates on Y, then the equations of Γ are

$$dy^p = F^p_i(x,y)\, dx^i \tag{2}$$

2010 *Mathematics Subject Classification*: primary 53C05; secondary 58A20, 58A32.
Key words and phrases: general connection, tangent valued form, functorial prolongation, Weil functor.
Received November 29, 2017. Editor J. Slovák.
DOI: 10.5817/AM2018-2-111

with arbitrary smooth functions F_i^p. Every vector field X on M defines the Γ-lift $\Gamma(X)\colon Y \to TY$, $\Gamma(X)(y) = \Gamma(y, X)$. Write $\pi_M\colon TM \to M$ for the bundle projection.

Equivalently, Γ can be interpreted as a section $Y \to J^1Y$ of the first jet prolongation J^1Y of Y. It is well known that $J^1Y \to Y$ is an affine bundle with associated vector bundle $VY \otimes T^*M$, where VY is the vertical tangent bundle of Y. For a section $s\colon M \to Y$, its absolute differential $\nabla_\Gamma s$ with respect to Γ is a section $\nabla_\Gamma s\colon M \to VY \otimes T^*M$ defined by

$$\nabla_\Gamma s(x) = j^1_x s - \Gamma\big(s(x)\big) \tag{3}$$

$x \in M$. Hence the coordinate form of (3) is

$$\frac{\partial s^p}{\partial x^i} - F_i^p\big(x, s(x)\big)\,. \tag{4}$$

The curvature $C\Gamma\colon Y \times_M \Lambda^2 T^*M \to VY$ can be characterized as the obstruction for lifting the bracket

$$(C\Gamma)(y, X_1, X_2) = \big[\Gamma(X_1), \Gamma(X_2)\big](y) - \Gamma\big([X_1, X_2]\big)(y)\,. \tag{5}$$

By direct evaluation, we find that (5) depends on the values of the vector fields X_1, X_2 at $p(y)$ only and the coordinate form of (5) is

$$2\Big(\frac{\partial F_i^p}{\partial x^j} + \frac{\partial F_i^p}{\partial y^q}\,F_j^q\Big)\frac{\partial}{\partial y^p} \otimes dx^i \wedge dx^j\,. \tag{6}$$

Using the flow prolongation of vector fields, we construct an induced connection $\mathcal{V}\Gamma\colon VY \times_M TM \to TVY$ on VY as follows, [6]. Consider the flow $\mathrm{Fl}_t^{\Gamma(X)}$ of the vector filed $\Gamma(X)$ and its vertical flow prolongation

$$\mathcal{V}\big(\Gamma(X)\big) = \frac{\partial}{\partial t}\Big|_0 V\big(\mathrm{Fl}_t^{\Gamma(X)}\big)\colon VY \to TVY\,. \tag{7}$$

Write $\eta^p = dy^p$ for the induced coordinates on VY. Then the coordinate form of (7) is

$$\begin{aligned} dy^p &= F_i^p(x, y)\,dx^i\,,\\ d\eta^p &= \frac{\partial F_i^p}{\partial y^q}\,\eta^q\,dx^i\,, \end{aligned} \tag{8}$$

that determines a general connection $\mathcal{V}\Gamma$ on $VY \to M$. The theoretical meaning of the vertical operator $\mathcal{V}$ is underlined by the following assertion, [6].

Proposition 1. *$\mathcal{V}$ is the only natural operator transforming general connections on $Y \to M$ into general connections on $VY \to M$.*

Consider a section $\varphi\colon Y \to VY \otimes \Lambda^k T^*M$ with the coordinate expression

$$\eta^p = \varphi^p_{i_1\dots i_k}(x, y)\,dx^{i_1} \wedge \dots \wedge dx^{i_k}\,.$$

According to [6], we construct its absolute exterior differential

$$d_{\mathcal{V}\Gamma}\varphi\colon Y \to VY \otimes \bigwedge^{k+1} T^*M$$

as follows. Take (at least locally) an auxiliary linear symmetric connection Λ on M. Then $\mathcal{V}\Gamma \otimes \overset{k}{\bigwedge} \Lambda^*$ is a connection on $VY \otimes \overset{k}{\bigwedge} T^*M \to Y$ and we can construct the absolute differential

$$\nabla_{\mathcal{V}\Gamma \otimes \bigwedge^k \Lambda^*} \varphi \colon Y \to V\big(VY \otimes \overset{k}{\bigwedge} T^*M\big) \otimes T^*M\,,$$

[6]. Applying antisymmetrization and natural identifications, we obtain a section $d_{\mathcal{V}\Gamma}\varphi \colon Y \to VY \otimes \Lambda^{k+1}T^*M$ independent of Λ with the coordinate expression

$$\eta^p = \Big(\frac{\partial \varphi^p_{i_1\dots i_k}}{\partial x^i} + \frac{\partial \varphi^p_{i_1\dots i_k}}{\partial y^q}\, F^q_i - \frac{\partial F^p_i}{\partial y^q}\, \varphi^q_{i_1\dots i_k}\Big)\, dx^i \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}\,. \tag{9}$$

In [6], we deduced by direct evaluation

Proposition 2 (Bianchi identity)**.** *We have*

$$d_{\mathcal{V}\Gamma} C\Gamma = 0\,. \tag{10}$$

2. Tangent valued forms

Mangiarotti and Modugno studied systematically the general connections by using the concept of tangent valued forms, [7]. A tangent valued k-form P on a manifold M is a section $P \colon M \to TM \otimes \Lambda^k T^*M$, that can be also interpreted as a map

$$P \colon TM \underbrace{\times_M \dots \times_M}_{k\text{-times}} TM \to TM\,. \tag{11}$$

If Q is another tangent valued l-form on M, Mangiarotti and Modugno defined a tangent valued $(k+l)$-form $[P,Q]$ on M by the formula

$$\begin{aligned}
&\big[P,Q\big](X_1 \dots, X_{k+l}) \\
&\quad = \frac{1}{k!l!} \sum_\sigma \overline{\sigma}\big[P(X_{\sigma_1}, \dots, X_{\sigma_k}), Q(X_{\sigma_{(k+1)}}, \dots, X_{\sigma_{(k+l)}})\big] \\
&\qquad + \frac{-1}{k!(l-1)!} \sum_\sigma \overline{\sigma} Q\big(\big[P(X_{\sigma_1}, \dots, X_{\sigma_k}), X_{\sigma_{(k+1)}}\big], X_{\sigma_{(k+2)}}, \dots\big) \\
&\qquad + \frac{(-1)^{kl}}{(k-1)!l!} \sum_\sigma \overline{\sigma} P\big(\big[Q(X_{\sigma_1}, \dots, X_{\sigma_l}), X_{\sigma_{(l+1)}}\big], X_{\sigma_{(l+2)}}, \dots\big) \\
&\qquad + \frac{(-1)^{k-1}}{(k-1)!(l-1)!2} \sum_\sigma \overline{\sigma} Q\big(P([X_{\sigma_1}, X_{\sigma_2}], X_{\sigma_3}, \dots), X_{\sigma_{(k+2)}}, \dots\big) \\
&\qquad + \frac{(-1)^{(k-1)l}}{(k-1)!(l-1)!2} \sum_\sigma \overline{\sigma} P\big(Q([X_{\sigma_1}, X_{\sigma_2}], X_{\sigma_3}, \dots), X_{\sigma_{(l+2)}}, \dots\big)
\end{aligned} \tag{12}$$

where $X_1, \dots, X_{k+l}$ are vector fields on M, the bracket on the right-hand side are the classical Lie bracket of vector fields, the summations are with respect to all permutations σ of $k+l$ letters and $\overline{\sigma}$ denotes the signum of σ. The tangent valued

0-forms are the vector fields and (12) reduces to the classical Lie bracket in the case $k = l = 0$.

Later it was clarified, [6], that (12) was introduced in a quite different situation by Frölicher-Nijenhuis, so that this bracket is related with their names today.

The identity of TM is a special tangent valued 1-form on M and we have

$$[\mathrm{id}_{TM}, P] = 0 \tag{13}$$

for every tangent valued form P. By [6],

$$[P, Q] = -(-1)^{kl}[Q, P] \tag{14}$$

and the graded Jacobi identity holds

$$\big[P_1, [P_2, P_3]\big] = \big[[P_1, P_2], P_3\big] + (-1)^{k_1 k_2}\big[P_2, [P_1, P_3]\big] \tag{15}$$

for tangent valued k_i-forms P_i, $i = 1, 2, 3$.

A general connection $\Gamma\colon Y \times_M TM \to TY$ defines a tangent valued 1-form ω_Γ on Y

$$\omega_\Gamma(Z) = \Gamma\big(y, Tp(Z)\big)\,, \quad Z \in T_yY\,. \tag{16}$$

Even $C\Gamma$ can be interpreted as a tangent valued 2-form C_Γ on Y,

$$C_\Gamma(Z_1, Z_2) = C\Gamma\big(y, Tp(Z_1), Tp(Z_2)\big)\,, \quad Z_1, Z_2 \in T_yY\,. \tag{17}$$

Proposition 3. *We have* $C_\Gamma = \frac{1}{2}[\omega_\Gamma, \omega_\Gamma]$.

Proof. This follows directly from Lemma 8.13 in [6]. □

Consider an arbitrary tangent valued 1-form ψ of Y. Put $P_1 = P_2 = P_3 = \psi$ into (14) and (15) This yields

$$\big[\psi, [\psi, \psi]\big] = 0\,.$$

If $\psi = \omega_\Gamma$, we obtain

Proposition 4. *We have* $[\omega_\Gamma, [\omega_\Gamma, \omega_\Gamma]] = 0$.

A simple evaluation shows that this relation coincides with the identity from Proposition 2. This gives a simple geometric proof of the Bianchi identity of a general connection Γ on Y.

3. Weilian prolongations

We recall that Weil algebra is a finite dimensional, commutative, associative and unital algebra of the form $A = \mathbb{R} \times N$, where N is the ideal of all nilpotent elements of A. Since A is finite dimensional, there exists an integer r such that $N^{r+1} = 0$. The smallest r with this property is called the order of A. On the other hand, the dimension wA of the vector space N/N^2 is the width of A, [8]. Using systematically our point of view, we say that a Weil algebra of width k and order r is a Weil (k, r)-algebra, [5].

The simpliest example of a Weil (k, r)-algebra is

$$\mathbb{D}^r_k = \mathbb{R}[x_1, \dots, x_k]\big/\langle x_1, \dots, x_k\rangle^{r+1} = J^r_0(\mathbb{R}^k, \mathbb{R})\,.$$

For $k = r = 1$, $\mathbb{D}^1_1 = \mathbb{D}$ is the algebra of Study numbers. In [3] we deduced

Lemma 1. *Every Weil (k,r)-algebra is a factor algebra of $\mathbb{D}^r_k$. If ϱ, $\sigma\colon \mathbb{D}^r_k \to A$ are two algebra epimorphisms, then there exists an algebra isomorphism $\chi\colon \mathbb{D}^r_k \to \mathbb{D}^r_k$ such that $\varrho = \sigma \circ \chi$.*

We are going to present the covariant approach to Weil functors, [5].

Definition 1. Two maps γ, $\delta\colon \mathbb{R}^k \to M$ determine the same A-velocity $j^A\gamma = j^A\delta$, if for every smooth function $\varphi\colon M \to \mathbb{R}$,

$$\varrho\big(j^r_0(\varphi\circ\gamma)\big) = \varrho\big(j^r_0(\varphi\circ\delta)\big)\,. \tag{18}$$

By Lemma 1, this is independent of the choice of ϱ. We say that

$$T^AM = \{j^A\gamma; \gamma\colon \mathbb{R}^k \to M\} \tag{19}$$

is the bundle of all A-velocities on M. For every smooth map $f\colon M \to N$, we define $T^Af\colon T^AM \to T^AM$ by

$$T^Af(j^A\gamma) = j^A(f\circ\gamma)\,. \tag{20}$$

Clearly, $T^A\mathbb{R} = A$.

We say that (19) and (20) represent the covariant approach to Weil functors. The following result is a fundamental assertion, see [6] or [5] for a survey.

Theorem. *The product preserving bundle functors on $\mathcal{M}f$ are in bijection with T^A. The natural transformations $T^{A_1} \to T^{A_2}$ are in bijection with the algebra homomorphisms $\mu\colon A_1 \to A_2$.*

We write $\mu_M\colon T^{A_1}M \to T^{A_2}M$ for the value of $\mu\colon A_1 \to A_2$ on M.

The iteration $T^{A_2}\circ T^{A_1}$ corresponds to the tensor product of A_1 and A_2. The algebra exchange homomorphism $\mathrm{ex}\colon A_1\otimes A_2 \to A_2\otimes A_1$ defines a natural exchange transformation $T^{A_2}T^{A_1} \to T^{A_1}T^{A_2}$. We have $T = T^{\mathbb{D}}$.

The canonical exchange $\varkappa^A_M\colon T^ATM \to TT^AM$ is called flow natural. Indeed, if Fl^X_t is the flow of a vector field $X\colon M \to TM$, then

$$\mathcal{T}^AX = \frac{\partial}{\partial t}\Big|_0 T^A(\mathrm{Fl}^X_t)\colon T^AM \to TT^AM$$

is the flow prolongation of X. It is related with the functorial prolongation $T^AX\colon T^AM \to T^ATM$ by

$$\mathcal{T}^AX = \varkappa^A_M \circ T^AX\,. \tag{21}$$

Consider a tangent valued k-form P on a manifold M

$$P\colon TM \times_M \cdots \times_M TM \to TM\,.$$

Applying functor T^A, we obtain

$$T^AP\colon T^ATP \times_M \cdots \times_M T^ATP \to T^ATP\,.$$

Using the flow natural exchange $\varkappa^A_M$, we construct

$$\mathcal{T}^AP = \varkappa^A_M \circ T^AP \circ \big((\varkappa^A_M)^{-1} \times \cdots \times (\varkappa^A_M)^{-1}\big)\,. \tag{22}$$

This is an antisymmetric tensor field of type $(1,k)$, so a tangent valued k-form on T^AM.

In [1], the following result is deduced.

Proposition 5. *The Frölicher-Nijenhuis bracket is preserved under $\mathcal{T}^A$, i.e. for every tangent valued k-form P and every tangent valued l-form Q on the same manifold M, we have*

$$\mathcal{T}^A([P,Q]) = [\mathcal{T}^A P, \mathcal{T}^A Q]\,. \tag{23}$$

Further, consider a tangent valued k-form P on a manifold M, a tangent valued k-form Q on a manifold N and a smooth map $f\colon M \to N$. We say that P and Q are f-related, if the following diagram commutes

$$\begin{array}{ccc} \Lambda^k TM & \xrightarrow{\quad P \quad} & TM \\ {\scriptstyle \Lambda^k Tf}\downarrow & & \downarrow{\scriptstyle Tf} \\ \Lambda^k TN & \xrightarrow{\quad Q \quad} & TN \end{array}$$

In [6], p. 74, one has deduced

Proposition 6. *Consider a smooth map $f\colon M \to N$. Let P_1, Q_1 or P_2, Q_2 be two f-related pairs of k-forms or l-forms, respectively. Then the Frölicher-Nijenhuis brackets $[P_1, Q_1]$ and $[P_2, Q_2]$ are also f-related.*

Consider a general connection Γ on Y in the lifting form $\Gamma\colon Y \times_M TM \to TY$. Applying T^A, $\varkappa^A_M$ and $\varkappa^A_Y$, [4, 5], we can construct the induced connection on $T^AY \to T^AM$

$$\mathcal{T}^A\Gamma\colon T^AY \times_{T^AM} TT^AM \to TT^AY\,. \tag{24}$$

Consider the connection form $\omega_\Gamma\colon TY \to TY$ of Γ. Then Proposition 5 and (24) imply

$$\mathcal{T}^A C_\Gamma = \frac{1}{2}\big[\mathcal{T}^A\omega_\Gamma, \mathcal{T}^A\omega_\Gamma\big]\,. \tag{25}$$

Hence the curvature of $\mathcal{T}^A\Gamma$ is the $\mathcal{T}^A$-prolongation of the curvature of Γ.

Further, the Bianchi identity of $\mathcal{T}^A\Gamma$ is the $\mathcal{T}^A$-prolongation of the Bianchi identity of Γ.

References

[1] Cabras, A., Kolář, I., *Prolongation of tangent valued forms to Weil bundles*, Arch. Math. (Brno) **31** (1995), 139–145.

[2] Ehresmann, C., *Oeuvres complètes et commentés*, Cahiers Topol. Géom. Diff. XXIV (Suppl. 1 et 2) (1983).

[3] Kolář, I., *Handbook of Global Analysis*, ch. Weil Bundles as Generalized Jet Spaces, pp. 625–665, Elsevier, Amsterdam, 2008.

[4] Kolář, I., *On the functorial prolongations of fiber bundles*, Miskolc Math. Notes **14** (2013), 423–431.

[5] Kolář, I., *Covariant Approach to Weil Bundles*, Folia, Masaryk University, Brno (2016).

[6] Kolář, I., Michor, P.W., Slovák, J., *Natural Operations in Differential Geometry*, Springer Verlag, 1993.

[7] Mangiarotti, L., Modugno, M., *Graded Lie algebras and connections on a fibered space*, J. Math. Pures Appl. (9) **63** (1984), 111–120.

[8] Weil, A., *Théorie des points proches sur les variétes différentielles*, Colloque de topol. et géom. diff., Strasbourg (1953), 111–117.

Institute of Mathematics and Statistics,
Masaryk University, Kotlářská 2,
611 37 Brno, Czech Republic
E-mail: kolar@math.muni.cz