ARCHIVUM MATHEMATICUM (BRNO)
Tomus 56 (2020), 225–247

# ON RIEMANN-POISSON LIE GROUPS

Brahim Alioune[a], Mohamed Boucetta[b], and Ahmed Sid'Ahmed Lessiad[c]

Abstract. A Riemann-Poisson Lie group is a Lie group endowed with a left invariant Riemannian metric and a left invariant Poisson tensor which are compatible in the sense introduced in [4]. We study these Lie groups and we give a characterization of their Lie algebras. We give also a way of building these Lie algebras and we give the list of such Lie algebras up to dimension 5.

## 1. Introduction

In this paper, we study Lie groups endowed with a left invariant Riemannian metric and a left invariant Poisson tensor satisfying a compatibility condition to be defined below. They constitute a subclass of the class of *Riemann-Poisson manifolds* introduced and studied by the second author (see [2, 3, 4, 5]).

Let $(M, \pi, \langle\ ,\ \rangle)$ be smooth manifold endowed with a Poisson tensor $\pi$ and a Riemannian metric $\langle\ ,\ \rangle$. We denote by $\langle\ ,\ \rangle^*$ the Euclidean product on $T^*M$ naturally associated to $\langle\ ,\ \rangle$. The Poisson tensor defines a Lie algebroid structure on $T^*M$ where the anchor map is the contraction $\#_\pi\colon T^*M \longrightarrow TM$ given by $\prec \beta, \#_\pi(\alpha) \succ = \pi(\alpha, \beta)$ and the Lie bracket on $\Omega^1(M)$ is the Koszul bracket given by

$$[\alpha,\beta]_\pi = \mathcal{L}_{\#_\pi(\alpha)}\beta - \mathcal{L}_{\#_\pi(\beta)}\alpha - d\pi(\alpha,\beta), \quad \alpha,\beta \in \Omega^1(M)\,. \tag{1}$$

This Lie algebroid structure and the metric $\langle\ ,\ \rangle^*$ define a contravariant connection $\mathcal{D}\colon \Omega^1(M)\times\Omega^1(M) \longrightarrow \Omega^1(M)$ by Koszul formula

$$\begin{aligned} 2\langle \mathcal{D}_\alpha\beta,\gamma\rangle^* &= \#_\pi(\alpha).\langle\beta,\gamma\rangle^* + \#_\pi(\beta).\langle\alpha,\gamma\rangle^* - \#_\pi(\gamma).\langle\alpha,\beta\rangle^* \\ &\quad + \langle[\alpha,\beta]_\pi,\gamma\rangle^* + \langle[\gamma,\alpha]_\pi,\beta\rangle^* + \langle[\gamma,\beta]_\pi,\alpha\rangle^*, \quad \alpha,\beta,\gamma\in\Omega^1(M)\,. \end{aligned} \tag{2}$$

This is the unique torsionless contravariant connection which is metric, i.e., for any $\alpha$, $\beta$, $\gamma \in \Omega^1(M)$,

$$\mathcal{D}_\alpha\beta - \mathcal{D}_\beta\alpha = [\alpha,\beta]_\pi \quad \text{and} \quad \#_\pi(\alpha).\langle\beta,\gamma\rangle^* = \langle\mathcal{D}_\alpha\beta,\gamma\rangle^* + \langle\beta,\mathcal{D}_\alpha\gamma\rangle^*\,.$$

The notion of contravariant connection was introduced by Vaisman in [13] and studied in more details by Fernandes in the context of Lie algebroids [8]. The

2020 *Mathematics Subject Classification*: primary 53A15; secondary 53D17, 22E05.
*Key words and phrases*: Lie group, Poisson manifolds, Riemannian metric.
Received August 22, 2019. Editor J. Slovák.
DOI: 10.5817/AM2020-4-225

connection $\mathcal{D}$ defined above is called *contravariant Levi-Civita connection* associated to the couple $(\pi, \langle\ ,\ \rangle)$ and it appeared first in [2].

The triple $(M, \pi, \langle\ ,\ \rangle)$ is called a *Riemannian-Poisson manifold* if $\mathcal{D}\pi = 0$, i.e., for any $\alpha, \beta, \gamma \in \Omega^1(M)$,

$$\mathcal{D}\pi(\alpha, \beta, \gamma) := \#_\pi(\alpha).\pi(\beta, \gamma) - \pi(\mathcal{D}_\alpha\beta, \gamma) + \pi(\beta, \mathcal{D}_\alpha\gamma) = 0\,. \tag{3}$$

This notion was introduced by the second author in [2]. Riemann-Poisson manifolds turned out to have interesting geometric properties (see[2, 3, 4, 5]). Let's mention some of them.

(1) The condition of compatibility (3) is weaker than the condition $\nabla\pi = 0$ where $\nabla$ is the Levi-Civita connection of $\langle\ ,\ \rangle$. Indeed, the condition (3) allows the Poisson tensor to have a variable rank. For instance, linear Poisson structures which are Riemann-Poisson exist and were characterized in [5]. Furthermore, let $(M, \langle\ ,\ \rangle)$ be a Riemannian manifold and $(X_1, \ldots, X_r)$ a family of commuting Killing vector fields. Put

$$\pi = \sum_{i,j} X_i \wedge X_j\,.$$

Then $(M, \pi, \langle\ ,\ \rangle)$ is a Riemann-Poisson manifold. This example illustrates also the weakness of the condition (3) and, more importantly, it is the local model of the geometry of noncommutative deformations studied by Hawkins (see [10, Theorem 6.6]).

(2) Riemann-Poisson manifolds can be thought of as a generalization of Kähler manifolds. Indeed, let $(M, \pi, \langle\ ,\ \rangle)$ be a Poisson manifold endowed with a Riemannian metric such that $\pi$ is invertible. Denote by $\omega$ the symplectic form inverse of $\pi$. Then $(M, \pi, \langle\ ,\ \rangle)$ is Riemann-Poisson manifold if and only if $\nabla\omega = 0$ where $\nabla$ is the Levi-Civita connection of $\langle\ ,\ \rangle$. In this case, if we define $A\colon TM \longrightarrow TM$ by $\omega(u, v) = \langle Au, v\rangle$ then $-A^2$ is symmetric definite positive and hence there exists a unique $Q\colon TM \longrightarrow TM$ symmetric definite positive such that $Q^2 = -A^2$. It follows that $J = AQ^{-1}$ satisfies $J^2 = -\mathrm{Id}_{TM}$, skew-symmetric with respect $\langle\ ,\ \rangle$ and $\nabla J = 0$. Hence $(M, J, \langle\ ,\ \rangle)$ is a Kähler manifold and its Kähler form $\omega_J(u, v) = \langle Ju, v\rangle$ is related to $\omega$ by the following formula:

$$\omega(u, v) = -\omega_J\left(\sqrt{-A^2}u, v\right), \quad u, v \in TM\,. \tag{4}$$

Having this construction in mind, we will call in this paper a Kähler manifold a triple $(M, \langle\ ,\ \rangle, \omega)$ where $\langle\ ,\ \rangle$ is a Riemannian metric and $\omega$ is a nondegenerate 2-form $\omega$ such that $\nabla\omega = 0$ where $\nabla$ is the Levi-Civita connection of $\langle\ ,\ \rangle$.

(3) The symplectic foliation of a Riemann-Poisson manifold when $\pi$ has a constant rank has an important property namely it is both a Riemannian foliation and a Kähler foliation.

Recall that a Riemannian foliation is a foliated manifold $(M, \mathcal{F})$ with a Riemannian metric $\langle\ ,\ \rangle$ such that the orthogonal distribution $T^\perp\mathcal{F}$ is totally geodesic.

Kähler foliations are a generalization of Kähler manifolds (see [6]) and, as for the notion of Kähler manifold, we call in this paper a Kähler foliation a foliated manifold $(M, \mathcal{F})$ endowed with a leafwise metric $\langle\ ,\ \rangle_{\mathcal{F}} \in \Gamma(\otimes^2 T^*\mathcal{F})$ and a nondegenerate leafwise differential 2-form $\omega_{\mathcal{F}} \in \Gamma(\otimes^2 T^*\mathcal{F})$ such any leaf with the restrictions of $\langle\ ,\ \rangle_{\mathcal{F}}$ and $\omega_{\mathcal{F}}$ is a Kähler manifold.

**Theorem 1.1** ([4]). *Let $(M, \langle\ ,\ \rangle, \pi)$ be a Riemann-Poisson manifold with $\pi$ of constant rank. Then its symplectic foliation is both a Riemannian and a Kähler foliation.*

Having in mind these properties particularly Theorem 1.1, it will be interesting to find large classes of examples of Riemann-Poisson manifolds. This paper will describe the rich collection of examples which are obtained by providing an arbitrary Lie group $G$ with a Riemannian metric $\langle\ ,\ \rangle$ and a Poisson tensor $\pi$ invariant under left translations and such that $(G, \langle\ ,\ \rangle, \pi)$ is Riemann-Poisson. We call $(G, \langle\ ,\ \rangle, \pi)$ a *Riemann-Poisson Lie group*. This class of examples can be enlarged substantially, with no extra work, as follows. If $(G, \langle\ ,\ \rangle, \pi)$ is a Riemann-Poisson Lie group and $\Gamma$ is any discrete subgroup of $G$ then $\Gamma\backslash G$ carries naturally a structure of Riemann-Poisson manifold.

The paper is organized as follows. In Section 2, we give the material needed in the paper and we describe the infinitesimal counterpart of Riemann-Poisson Lie groups, namely, Riemann-Poisson Lie algebras. In Section 3, we prove our main result which gives an useful description of Riemann-Poisson Lie algebras (see Theorem 3.1). We use this theorem in Section 4 to derive a method for building Riemann-Poisson Lie algebras. We explicit this method by giving the list of Riemann-Poisson Lie algebras up to dimension 5.

## 2. Riemann-Poisson Lie groups and their infinitesimal characterization

Let $G$ be a Lie group and $(\mathfrak{g} = T_eG, [\ ,\ ])$ its Lie algebra.

(1) A left invariant Poisson tensor $\pi$ on $G$ is entirely determined by

$$\pi(\alpha, \beta)(a) = r(\mathrm{L}_a^*\alpha, \mathrm{L}_a^*\beta)\,,$$

where $a \in G$, $\alpha$, $\beta \in T_a^*G$, $\mathrm{L}_a$ is the left multiplication by $a$ and $r \in \wedge^2\mathfrak{g}$ satisfies the classical Yang-Baxter equation

$$[r, r] = 0\,, \tag{5}$$

where $[r, r] \in \wedge^3\mathfrak{g}$ is given by

$$\begin{aligned}[r, r](\alpha, \beta, \gamma) :=& \prec \alpha, [r_\#(\beta), r_\#(\gamma)] \succ + \prec \beta, [r_\#(\gamma), r_\#(\alpha)] \succ \\ &+ \prec \gamma, [r_\#(\alpha), r_\#(\beta)] \succ\,, \quad \alpha, \beta, \gamma \in \mathfrak{g}^*,\end{aligned} \tag{6}$$

and $r_\#\colon \mathfrak{g}^* \longrightarrow \mathfrak{g}$ is the contraction associated to $r$. In this case, the Koszul bracket (1) when restricted to left invariant differential 1-forms induces a Lie bracket on $\mathfrak{g}^*$ given by

$$[\alpha, \beta]_r = \mathrm{ad}^*_{r_\#(\alpha)}\beta - \mathrm{ad}^*_{r_\#(\beta)}\alpha\,, \quad \alpha, \beta \in \mathfrak{g}^*, \tag{7}$$

where $\prec \mathrm{ad}^*_u\alpha, v \succ = - \prec \alpha, [u,v] \succ$. Moreover, $r_\#$ becomes a morphism of Lie algebras, i.e.,

$$r_\#([\alpha,\beta]_r) = [r_\#(\alpha), r_\#(\beta)]\,, \quad \alpha,\beta \in \mathfrak{g}^*. \tag{8}$$

(2) A let invariant Riemannian metric $\langle\ ,\ \rangle$ on $G$ is entirely determined by

$$\langle u, v\rangle(a) = \rho(T_a\mathrm{L}_{a^{-1}}u, T_a\mathrm{L}_{a^{-1}}v),$$

where $a \in G$, $u$, $v \in T_aG$ and $\rho$ is a scalar product on $\mathfrak{g}$. The Levi-Civita connection of $\langle\ ,\ \rangle$ is left invariant and induces a product $A\colon \mathfrak{g}\times\mathfrak{g} \longrightarrow \mathfrak{g}$ given by

$$2\varrho(A_uv, w) = \varrho([u,v],w) + \varrho([w,u],v) + \varrho([w,v],u)\,, \quad u,v,w \in \mathfrak{g}\,. \tag{9}$$

It is the unique product on $\mathfrak{g}$ satisfying

$$A_uv - A_vu = [u,v] \quad \text{and} \quad \varrho(A_uv,w) + \varrho(v, A_uw) = 0\,,$$

for any $u$, $v$, $w \in \mathfrak{g}$. We call $A$ the *Levi-Civita product* associated to $(\mathfrak{g}, [\ ,\ ], \rho)$.

(3) Let $(G, \langle\ ,\ \rangle, \Omega)$ be a Lie group endowed with a left invariant Riemannian metric and a nondegenerate left invariant 2-form. Then $(G, \langle\ ,\ \rangle, \Omega)$ is a Kähler manifold if and only if, for any $u, v, w \in \mathfrak{g}$,

$$\omega(A_uv, w) + \omega(u, A_uv) = 0\,, \tag{10}$$

where $\omega = \Omega(e)$, $\rho = \langle\ ,\ \rangle(e)$ and $A$ is the Levi-Civita product of $(\mathfrak{g}, [\ ,\ ], \rho)$. In this case we call $(\mathfrak{g}, [\ ,\ ], \rho, \omega)$ a Kähler Lie algebra.

As all the left invariant structures on Lie groups, Riemann-Poison Lie groups can be characterized at the level of their Lie algebras.

**Proposition 2.1.** *Let $(G, \pi, \langle\ ,\ \rangle)$ be a Lie group endowed with a left invariant bivector field and a left invariant metric and $(\mathfrak{g}, [\ ,\ ])$ its Lie algebra. Put $r = \pi(e) \in \wedge^2\mathfrak{g}$, $\varrho = \langle\ ,\ \rangle_e$ and $\varrho^*$ the associated Euclidean product on $\mathfrak{g}^*$. Then $(G, \pi, \langle\ ,\ \rangle)$ is a Riemann-Poisson Lie group if and only if*

(i) $[r,r] = 0$,

(ii) *for any $\alpha$, $\beta$, $\gamma \in \mathfrak{g}^*$, $r(A_\alpha\beta, \gamma) + r(\beta, A_\alpha\gamma) = 0$,*

*where $A$ is the Levi-Civita product associated to $(\mathfrak{g}^*, [\ ,\ ]_r, \varrho^*)$.*

**Proof.** For any $u \in \mathfrak{g}$ and $\alpha \in \mathfrak{g}^*$, we denote by $u^\ell$ and $\alpha^\ell$, respectively, the left invariant vector field and the left invariant differential 1-form on $G$ given by

$$u^\ell(a) = T_e\mathrm{L}_a(u) \quad \text{and} \quad \alpha^\ell(a) = T^*_a\mathrm{L}_{a^{-1}}(\alpha)\,, \quad a \in G,\ \mathrm{L}_a(b) = ab\,.$$

Since $\pi$ and $\langle\ ,\ \rangle$ are left invariant, one can see easily from (1) and (2) that we have, for any $\alpha, \beta, \gamma \in \mathfrak{g}^*$,

$$\begin{cases} [\pi,\pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = [r,r](\alpha,\beta,\gamma),\ \#_\pi(\alpha^\ell) = (r_\#(\alpha))^\ell,\ \mathcal{L}_{\#_\pi(\alpha^\ell)}\beta^\ell = \left(\mathrm{ad}^*_{r_\#(\alpha)}\beta\right)^\ell, \\ [\alpha^\ell, \beta^\ell]_\pi = ([\alpha,\beta]_r)^\ell\,,\ \mathcal{D}_{\alpha^\ell}\beta^\ell = (A_\alpha\beta)^\ell\,. \end{cases}$$

The proposition follows from these formulas, (3) and the fact that $(G, \pi, \langle\ ,\ \rangle)$ is a Riemann-Poisson Lie group if and only if, for any $\alpha$, $\beta$, $\gamma \in \mathfrak{g}^*$,

$$[\pi, \pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0 \quad \text{and} \quad \mathcal{D}\pi(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0\,. \qquad \square$$

Conversely, given a triple $(\mathfrak{g}, r, \varrho)$ where $\mathfrak{g}$ is a real Lie algebra, $r \in \wedge^2\mathfrak{g}$ and $\varrho$ a Euclidean product on $\mathfrak{g}$ satisfying the conditions $(i)$ and $(ii)$ in Proposition 2.1 then, for any Lie group $G$ whose Lie algebra is $\mathfrak{g}$, if $\pi$ and $\langle\ ,\ \rangle$ are the left invariant bivector field and the left invariant metric associated to $(r, \varrho)$ then $(G, \pi, \langle\ ,\ \rangle)$ is a Riemann-Poisson Lie group.

**Definition 2.1.** A *Riemann-Poisson Lie algebra* is a triple $(\mathfrak{g}, r, \varrho)$ where $\mathfrak{g}$ is a real Lie algebra, $r \in \wedge^2\mathfrak{g}$ and $\varrho$ a Euclidean product on $\mathfrak{g}$ satisfying the conditions (i) and (ii) in Proposition 2.1.

To end this section, we give another characterization of the solutions of the classical Yang-Baxter equation (5) which will be useful later.

We observe that $r \in \wedge^2\mathfrak{g}$ is equivalent to the data of a vector subspace $S \subset \mathfrak{g}$ and a nondegenerate 2-form $\omega_r \in \wedge^2 S^*$.

Indeed, for $r \in \wedge^2\mathfrak{g}$, we put $S = \mathrm{Im} r_\#$ and $\omega_r(u, v) = r(r_\#^{-1}(u), r_\#^{-1}(v))$ where $u$, $v \in S$ and $r_\#^{-1}(u)$ is any antecedent of $u$ by $r_\#$.

Conversely, let $(S, \omega)$ be a vector subspace of $\mathfrak{g}$ with a non-degenerate 2-form. The 2-form $\omega$ defines an isomorphism $\omega^b \colon S \longrightarrow S^*$ by $\omega^b(u) = \omega(u, .)$, we denote by $\# \colon S^* \longrightarrow S$ its inverse and we put $r_\# = \# \circ i^*$ where $i^* \colon \mathfrak{g}^* \longrightarrow S^*$ is the dual of the inclusion $i \colon S \hookrightarrow \mathfrak{g}$.

With this observation in mind, the following proposition gives another description of the solutions of the Yang-Baxter equation.

**Proposition 2.2.** *Let $r \in \wedge^2\mathfrak{g}$ and $(S, \omega_r)$ its associated vector subspace. The following assertions are equivalent:*

(1) $[r, r] = 0$.

(2) *$S$ is a subalgebra of $\mathfrak{g}$ and*

$$\delta\omega_r(u, v, w) := \omega_r(u, [v, w]) + \omega_r(v, [w, u]) + \omega_r(w, [u, v]) = 0$$

*for any $u$, $v$, $w \in S$.*

**Proof.** The proposition follows from the following formulas:

$$\prec \gamma, r_\#([\alpha, \beta]_r) - [r_\#(\alpha), r_\#(\beta)] \succ = -[r, r](\alpha, \beta, \gamma)\,, \qquad \alpha, \beta, \gamma \in \mathfrak{g}^*$$

and, if $S$ is a subalgebra,

$$[r, r](\alpha, \beta, \gamma) = -\delta\omega_r(r_\#(\alpha), r_\#(\beta), r_\#(\gamma))\,. \quad \square$$

This proposition shows that there is a correspondence between the set of solutions of the Yang-Baxter equation the set of symplectic subalgebras of $\mathfrak{g}$. We recall that a symplectic algebra is a Lie algebra $S$ endowed with a non-degenerate 2-form $\omega$ such that $\delta\omega = 0$.

## 3. A characterization of Riemann-Poisson Lie algebras

In this section, we combine Propositions 2.1 and 2.2 to establish a characterization of Riemann-Poisson Lie algebras which will be used later to build such Lie algebras. We establish first an intermediary result.

**Proposition 3.1.** *Let $(\mathfrak{g}, r, \varrho)$ be a Lie algebra endowed with $r \in \wedge^2\mathfrak{g}$ and a Euclidean product $\varrho$. Denote by $\mathcal{I} = \ker r_\#$, $\mathcal{I}^\perp$ its orthogonal with respect to $\varrho^*$ and $A$ the Levi-Civita product associated to $(\mathfrak{g}^*, [\ ,\ ]_r, \varrho^*)$. Then $(\mathfrak{g}, r, \varrho)$ is a Riemann-Poisson Lie algebra if and only if:*

($c_1$) $[r, r] = 0$.

($c_2$) *For all $\alpha \in \mathcal{I}$, $A_\alpha = 0$.*

($c_3$) *For all $\alpha$, $\beta$, $\gamma \in \mathcal{I}^\perp$, $A_\alpha\beta \in \mathcal{I}^\perp$ and*

$$r(A_\alpha\beta, \gamma) + r(\beta, A_\alpha\gamma) = 0\,.$$

**Proof.** By using the splitting $\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^\perp$, on can see that the conditions $(i)$ and $(ii)$ in Proposition 2.1 are equivalent to

$$\begin{cases} [r, r] = 0\,, \\ r(A_\alpha\beta, \gamma) = 0\,,\ \alpha \in \mathcal{I}\,,\ \beta \in \mathcal{I}\,,\ \gamma \in \mathcal{I}^\perp, \\ r(A_\alpha\beta, \gamma) + r(\beta, A_\alpha\gamma) = 0\,,\ \alpha \in \mathcal{I}\,,\ \beta \in \mathcal{I}^\perp\,,\ \gamma \in \mathcal{I}^\perp, \\ r(A_\alpha\beta, \gamma) = 0\,,\ \alpha \in \mathcal{I}^\perp\,,\ \beta \in \mathcal{I}\,,\ \gamma \in \mathcal{I}^\perp, \\ r(A_\alpha\beta, \gamma) + r(\beta, A_\alpha\gamma) = 0\,,\ \alpha \in \mathcal{I}^\perp,\ \beta \in \mathcal{I}^\perp,\ \gamma \in \mathcal{I}^\perp. \end{cases} \tag{11}$$

Suppose that the conditions ($c_1$)-($c_3$) hold. Then for any $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^\perp$, $A_\beta\alpha = [\beta, \alpha]_r$ and hence $r_\#(A_\beta\alpha) = [r_\#(\beta), r_\#(\alpha)] = 0$ and hence the equations in (11) holds.

Conversely, suppose that (11) holds. Then ($c_1$) holds obviously.

For any $\alpha, \beta \in \mathcal{I}$, the second equation in (11) is equivalent to $A_\alpha\beta \in \mathcal{I}$ and we have from (7) and (9) $[\alpha, \beta]_r = 0$ and $A_\alpha\beta \in \mathcal{I}^\perp$. Thus $A_\alpha\beta = 0$.

Take now $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^\perp$. For any $\gamma \in \mathcal{I}$, $\varrho^*(A_\alpha\beta, \gamma) = -\varrho^*(\beta, A_\alpha\gamma) = 0$ and hence $A_\alpha\beta \in \mathcal{I}^\perp$. On the other hand,

$$r_\#([\alpha, \beta]_r) = r_\#(A_\alpha\beta) - r_\#(A_\beta\alpha) \overset{(8)}{=} [r_\#(\alpha), r_\#(\beta)] = 0\,.$$

So, for any $\gamma \in \mathcal{I}^\perp$,

$$\prec \gamma, r_\#(A_\alpha\beta) \succ = \prec \gamma, r_\#(A_\beta\alpha) \succ = r(A_\beta\alpha, \gamma) \overset{(11)}{=} 0\,.$$

This shows that $A_\alpha\beta \in \mathcal{I}$ and hence $A_\alpha\beta = 0$. Finally, ($c_2$) is true. Now, for any $\alpha \in \mathcal{I}^\perp$, the fourth equation in (11) implies that $A_\alpha$ leaves invariant $\mathcal{I}$ and since it is skew-symmetric it leaves invariant $\mathcal{I}^\perp$ and ($c_3$) follows. This completes the proof. □

**Proposition 3.2.** *Let $(\mathfrak{g}, \varrho, r)$ be a Lie algebra endowed with a solution of classical Yang-Baxter equation and a bi-invariant Euclidean product, i.e.,*

$$\varrho(\mathrm{ad}_u v, w) + \varrho(v, \mathrm{ad}_u w) = 0\,, \quad u, v, w \in \mathfrak{g}\,.$$

*Then* $(\mathfrak{g}, \varrho, r)$ *is Riemann-Poisson Lie algebra if and only if* $\mathrm{Im} r_\#$ *is an abelian subalgebra.*

**Proof.** Since $\varrho$ is bi-invariant, one can see easily that for any $u \in \mathfrak{g}$, $\mathrm{ad}_u^*$ is skew-symmetric with respect to $\varrho^*$ and hence the Levi-Civita product $A$ associated to $(\mathfrak{g}^*, [\ ,\ ]_r, \varrho^*)$ is given by $A_\alpha \beta = \mathrm{ad}^*_{r_\#(\alpha)}\beta$. So, $(\mathfrak{g}, \varrho, r)$ is Riemann-Poisson Lie algebra if and only if, for any $\alpha, \beta, \gamma \in \mathfrak{g}^*$,

$$\begin{aligned} 0 &= r(\mathrm{ad}^*_{r_\#(\alpha)}\beta, \gamma) + r(\beta, \mathrm{ad}^*_{r_\#(\alpha)}\gamma) \\ &= \prec \beta, [r_\#(\alpha), r_\#(\gamma)] \succ - \prec \gamma, [r_\#(\alpha), r_\#(\beta)] \succ \\ &\overset{(5)}{=} \prec \alpha, [r_\#(\beta), r_\#(\gamma)] \succ \end{aligned}$$

and the result follows. □

Let $(\mathfrak{g}, [\ ,\ ])$ be a Lie algebra, $r \in \wedge^2\mathfrak{g}$ and $\varrho$ a Euclidean product on $\mathfrak{g}$. Denote by $(S, \omega_r)$ the symplectic vector subspace associated to $r$ and by $\#\colon \mathfrak{g}^* \longrightarrow \mathfrak{g}$ the isomorphism given by $\varrho$. Note that the Euclidean product on $\mathfrak{g}^*$ is given by $\varrho^*(\alpha, \beta) = \varrho(\#(\alpha), \#(\beta))$. We have

$$\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^\perp \quad \text{and} \quad \mathfrak{g} = S \oplus S^\perp\,,$$

where $\mathcal{I} = \ker r_\#$. Moreover, $r_\#\colon \mathcal{I}^\perp \longrightarrow S$ is an isomorphism, we denote by $\tau\colon S \longrightarrow \mathcal{I}^\perp$ its inverse. From the relation

$$\varrho\big(\#(\alpha), r_\#(\beta)\big) = \prec \alpha\,,\ r_\#(\beta) \succ = r(\beta, \alpha)\,,$$

we deduce that $\#\colon \mathcal{I} \longrightarrow S^\perp$ is an isomorphism and hence $\#\colon \mathcal{I}^\perp \longrightarrow S$ is also an isomorphism.

Consider the isomorphism $J\colon S \longrightarrow S$ linking $\omega_r$ to $\varrho_{|S}$, i.e.,

$$\omega_r(u, v) = \rho(Ju, v)\,, \quad u, v \in S\,.$$

On can see easily that $J = -\# \circ \tau$.

**Theorem 3.1.** *With the notations above,* $(\mathfrak{g}, r, \varrho)$ *is a Riemann-Poisson Lie algebra if and only if the following conditions hold:*

(1) $(S, \varrho_{|S}, \omega_r)$ *is a Kähler Lie subalgebra, i.e., for all* $s_1$, $s_2$, $s_3 \in S$,

$$\omega_r(\nabla_{s_1} s_2, s_3) + \omega_r(s_2, \nabla_{s_1} s_3) = 0\,, \tag{12}$$

*where* $\nabla$ *is the Levi-Civita product associated to* $(S, [\ ,\ ], \varrho_{|S})$.

(2) *for all* $s \in S$ *and all* $u, v \in S^\perp$,

$$\varrho(\phi_S(s)(u), v) + \varrho(u, \phi_S(s)(v)) = 0\,, \tag{13}$$

*where* $\phi_S\colon S \longrightarrow \mathrm{End}(S^\perp)$, $u \mapsto \mathrm{pr}_{S^\perp} \circ \mathrm{ad}_u$ *and* $\mathrm{pr}_{S^\perp}\colon \mathfrak{g} \longrightarrow S^\perp$ *is the orthogonal projection.*

(3) *For all* $s_1$, $s_2 \in S$ *and all* $u \in S^\perp$,

$$\omega_r(\phi_{S^\perp}(u)(s_1), s_2) + \omega_r(s_1, \phi_{S^\perp}(u)(s_1)) = 0\,, \tag{14}$$

*where* $\phi_{S^\perp} : S^\perp \longrightarrow \mathrm{End}(S)$, $u \mapsto \mathrm{pr}_S \circ \mathrm{ad}_u$ *and* $\mathrm{pr}_S\colon \mathfrak{g} \longrightarrow S$ *is the orthogonal projection.*

**Proof.** Suppose first that $(\mathfrak{g}, r, \varrho)$ is a Riemann-Poisson Lie algebra. According to Propositions 3.1 and 2.2, this is equivalent to

$$\begin{cases} (S, \omega_r) \quad \text{is a symplectic subalgebra,} \\ \forall \alpha \in \mathcal{I},\ A_\alpha = 0\,, \\ \forall\ \alpha, \beta, \gamma \in \mathcal{I}^\perp,\ A_\alpha\beta \in \mathcal{I}^\perp \quad \text{and} \quad r(A_\alpha\beta, \gamma) + r(\beta, A_\alpha\gamma) = 0\,, \end{cases} \tag{15}$$

where $A$ is the Levi-Civita product of $(\mathfrak{g}^*, [\ ,\ ]_r, \varrho^*)$.

For $\alpha, \beta \in \mathcal{I}$ and $\gamma \in \mathcal{I}^\perp$,

$$\begin{aligned} 2\varrho^*(A_\alpha\beta, \gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= \varrho^*(\mathrm{ad}^*_{r_\#(\gamma)}\beta, \alpha) + \varrho^*(\mathrm{ad}^*_{r_\#(\gamma)}\alpha, \beta) \\ &= -\prec \beta, [r_\#(\gamma), \#(\alpha)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) - \varrho(\#(\alpha), [r_\#(\gamma), \#(\beta)])\,. \end{aligned} \tag{16}$$

Since $\#\colon \mathcal{I} \longrightarrow S^\perp$ and $r_\#\colon \mathcal{I}^\perp \longrightarrow S$ are isomorphisms, we deduce from (16) that $A_\alpha\beta = 0$ for any $\alpha,\ \beta \in \mathcal{I}$ is equivalent to (13).

For $\alpha \in \mathcal{I}$ and $\beta,\ \gamma \in \mathcal{I}^\perp$,

$$\begin{aligned} 2\varrho^*(A_\alpha\beta,\gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= -\varrho^*(\mathrm{ad}^*_{r_\#(\beta)}\alpha, \gamma) - \varrho^*(\mathrm{ad}^*_{r_\#(\beta)}\gamma, \alpha) + \varrho^*(\mathrm{ad}^*_{r_\#(\gamma)}\beta, \alpha) + \varrho^*(\mathrm{ad}^*_{r_\#(\gamma)}\alpha, \beta) \\ &= \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ + \prec \gamma, [r_\#(\beta), \#(\alpha)] \succ - \prec \beta, [r_\#(\gamma), \#(\alpha)] \succ \\ &\quad - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = \varrho(\#(\gamma), [r_\#(\beta), \#(\alpha)]) \\ &\quad - \varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(J \circ r_\#(\gamma), [r_\#(\beta), \#(\alpha)]) + \varrho(J \circ r_\#(\beta), [r_\#(\gamma), \#(\alpha)]) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\omega_r(r_\#(\gamma), \mathrm{pr}_S([r_\#(\beta), \#(\alpha)])) - \omega_r(\mathrm{pr}_S([r_\#(\gamma), \#(\alpha)]), r_\#(\beta)) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ\ . \end{aligned} \tag{17}$$

Now, $\#(\beta), \#(\gamma) \in S$ and $r_\#(\beta), r_\#(\gamma) \in S$ and since $S$ is a subalgebra we deduce that $[r_\#(\beta), \#(\gamma)], [r_\#(\gamma), \#(\beta)] \in S$ and hence

$$\prec \alpha, [r_\#(\beta), \#(\gamma)] \succ = \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = 0\,.$$

We have also $\#\colon \mathcal{I} \longrightarrow S^\perp$ and $r_\#\colon \mathcal{I}^\perp \longrightarrow S$ are isomorphisms so that, by virtue of (17), $A_\alpha\beta = 0$ for any $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^\perp$ is equivalent to (14).

On the other hand, for any $\alpha, \beta, \gamma \in \mathcal{I}^\perp$, since $\# = -J \circ r_\#$, the relation

$$2\varrho^*(A_\alpha\beta, \gamma) = \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta)$$

can be written

$$\begin{aligned} 2\varrho\big(J \circ r_\#(A_\alpha\beta), J \circ r_\#(\gamma)\big) &= \varrho\big(J \circ r_\#([\alpha, \beta]_r), J \circ r_\#(\gamma)\big) \\ &\quad + \varrho\big(J \circ r_\#([\gamma, \beta]_r), J \circ r_\#(\alpha)\big) \\ &\quad + \varrho\big(J \circ r_\#([\gamma, \alpha]_r), J \circ r_\#(\beta)\big)\,. \end{aligned}$$

But $r_\#([\alpha,\beta]_r) = [r_\#(\alpha), r_\#(\beta)]$ and hence

$$2\langle r_\#(A_\alpha\beta), r_\#(\gamma)\rangle_J = \langle [r_\#(\alpha), r_\#(\beta)], r_\#(\gamma)\rangle_J + \langle [r_\#(\mathfrak{g}), r_\#(\beta)], r_\#(\alpha)\rangle_J + \langle [r_\#(\gamma), r_\#(\alpha)], r_\#(\beta)\rangle_J ,$$

where $\langle u, v\rangle_J = \varrho(Ju, Jv)$. This shows that $r_\#(A_\alpha\beta) = \nabla_{r_\#(\alpha)} r_\#(\beta)$ where $\nabla$ is the Levi-Civita product of $(S, [\ ,\ ], \langle\ ,\ \rangle_J)$ and the third relation in (15) is equivalent to

$$\omega_r(\nabla_u v, w) + \omega_r(v, \nabla_u w) = 0 , \quad u, v, w \in S .$$

This is equivalent to $\nabla_u Jv = J\nabla_u v$. Let us show that $\nabla$ is actually the Levi-Civita product of $(S, [\ ,\ ], \varrho)$. Indeed, for any $u$, $v$, $w \in S$, $\nabla_u v - \nabla_v u = [u, v]$ and

$$\begin{aligned}\varrho(\nabla_u v, w) + \varrho(\nabla_u w, v) &= \langle J^{-1}\nabla_u v, J^{-1}w\rangle_J + \langle J^{-1}\nabla_u w, J^{-1}v\rangle_J \\ &= \langle \nabla_u J^{-1}v, J^{-1}w\rangle_J + \langle \nabla_u J^{-1}w, J^{-1}v\rangle_J \\ &= 0 .\end{aligned}$$

So we have shown the direct part of the theorem. The converse can be deduced easily from the relations we established in the proof of the direct part. □

**Example 1.** Let $G$ be a compact connected Lie group, $\mathfrak{g}$ its Lie algebra and $T$ an even dimensional torus of $G$. Choose a bi-invariant Riemannian metric $\langle\ ,\ \rangle$ on $G$, a nondegenerate $\omega \in \wedge^2 S^*$ where $S$ is the Lie algebra of $T$ and put $\varrho = \langle\ ,\ \rangle(e)$. Let $r \in \wedge^2\mathfrak{g}$ be the solution of the classical Yang-Baxter associated to $(S, \omega)$. By using either Proposition 3.2 or Theorem 3.1, one can see easily that $(\mathfrak{g}, \varrho, r)$ is a Riemann-Poisson Lie algebra and hence $(G, \langle\ ,\ \rangle, \pi)$ is a Riemann-Poisson Lie group where $\pi$ is the left invariant Poisson tensor associated to $r$. According to Theorem 1.1, the orbits of the right action of $T$ on $G$ defines a Riemannian and Kähler foliation. For instance, $G = \mathrm{SO}(2n)$, $T = \mathrm{Diagonal}(D_1, \ldots, D_n)$ where $D_i = \begin{pmatrix} \cos(\theta_i) & \sin(\theta_i) \\ -\sin(\theta_i) & \cos(\theta_i)\end{pmatrix}$ and $\langle\ ,\ \rangle = -K$ where $K$ is the Killing form.

## 4. Construction of Riemann-Poisson Lie algebras

In this section, we give a general method for building Riemann-Poisson Lie algebras and we use it to give all Riemann-Poisson Lie algebras up to dimension 5.

According to Theorem 3.1, to build Riemann-Poisson Lie algebras one needs to solve the following problem.

**Problem 1.** We look for:

(1) A Kähler Lie algebra $(\mathfrak{h}, [\ ,\ ]_\mathfrak{h}, \varrho_\mathfrak{h}, \omega)$,

(2) a Euclidean vector space $(\mathfrak{p}, \varrho_\mathfrak{p})$,

(3) a bilinear skew-symmetric map $[\ ,\ ]_\mathfrak{p} : \mathfrak{p}\times\mathfrak{p} \longrightarrow \mathfrak{p}$,

(4) a bilinear skew-symmetric map $\mu : \mathfrak{p}\times\mathfrak{p} \longrightarrow \mathfrak{h}$,

(5) two linear maps $\phi_\mathfrak{p} : \mathfrak{p} \longrightarrow \mathrm{sp}(\mathfrak{h}, \omega)$ and $\phi_\mathfrak{h} : \mathfrak{h} \longrightarrow \mathrm{so}(\mathfrak{p})$ where $\mathrm{sp}(\mathfrak{h}, \omega) = \{J : \mathfrak{h} \longrightarrow \mathfrak{h}, J^\omega + J = 0\}$ and $\mathrm{so}(\mathfrak{p}) = \{A : \mathfrak{p} \longrightarrow \mathfrak{p}, A^* + A = 0\}$, $J^\omega$ is the adjoint with respect to $\omega$ and $A^*$ is the adjoint with respect to $\varrho_\mathfrak{p}$,

such that the bracket $[\ ,\ ]$ on $\mathfrak{g}=\mathfrak{h}\oplus\mathfrak{p}$ given, for any $a,\ b\in\mathfrak{p}$ and $u,\ v\in\mathfrak{h}$, by

$$[u,v]=[u,v]_{\mathfrak{h}},\ [a,b]=\mu(a,b)+[a,b]_{\mathfrak{p}},\ [a,u]=-[u,a]=\phi_{\mathfrak{p}}(a)(u)-\phi_{\mathfrak{h}}(u)(a) \tag{18}$$

is a Lie bracket.

In this case, $(\mathfrak{g},[\ ,\ ])$ endowed with $r\in\wedge^2\mathfrak{g}$ associated to $(\mathfrak{h},\omega)$ and the Euclidean product $\varrho=\varrho_{\mathfrak{h}}\oplus\varrho_{\mathfrak{p}}$ becomes, by virtue of Theorem 3.1, a Riemann-Poisson Lie algebra.

**Proposition 4.1.** *With the data and notations of Problem 1, the bracket given by* (18) *is a Lie bracket if and only if, for any $u,\ v\in\mathfrak{h}$ and $a,\ b,\ c\in\mathfrak{p}$,*

(19)

$$\begin{cases}
\phi_{\mathfrak{p}}(a)([u,v]_{\mathfrak{h}})=[u,\phi_{\mathfrak{p}}(a)(v)]_{\mathfrak{h}}+[\phi_{\mathfrak{p}}(a)(u),v]_{\mathfrak{h}}+\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(v)(a))(u)-\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(u)(a))(v)\,,\\
\phi_{\mathfrak{h}}(u)([a,b]_{\mathfrak{p}})=[a,\phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}}+[\phi_{\mathfrak{h}}(u)(a),b]_{\mathfrak{p}}+\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a)-\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b)\,,\\
\phi_{\mathfrak{h}}([u,v]_{\mathfrak{h}})=[\phi_{\mathfrak{h}}(u),\phi_{\mathfrak{h}}(v)]\,,\\
\phi_{\mathfrak{p}}([a,b]_{\mathfrak{p}})(u)=[\phi_{\mathfrak{p}}(a),\phi_{\mathfrak{p}}(b)](u)+[u,\mu(a,b)]_{\mathfrak{h}}-\mu(a,\phi_{\mathfrak{h}}(u)(b))-\mu(\phi_{\mathfrak{h}}(u)(a),b)\,,\\
\oint[a,[b,c]_{\mathfrak{p}}]_{\mathfrak{p}}=\oint\phi_{\mathfrak{h}}(\mu(b,c))(a)\,,\\
\oint\phi_{\mathfrak{p}}(a)(\mu(b,c))=\oint\mu([b,c]_{\mathfrak{p}},a)\,,
\end{cases}$$

*where $\oint$ stands for the circular permutation.*

**Proof.** The equations follow from the Jacobi identity applied to $(a,u,v)$, $(a,b,u)$ and $(a,b,c)$. □

We tackle now the task of determining the list of all Riemann-Poisson Lie algebras up to dimension 5. For this purpose, we need to solve Problem 1 in the following four cases: $(a)$ $\dim\mathfrak{p}=1$, $(b)$ $\dim\mathfrak{h}=2$ and $\mathfrak{h}$ non abelian, $(c)$ $\dim\mathfrak{h}=\dim\mathfrak{p}=2$ and $\mathfrak{h}$ abelian, $(d)$ $\dim\mathfrak{h}=2$, $\dim\mathfrak{p}=3$ and $\mathfrak{h}$ abelian.

It is easy to find the solutions of Problem 1 when $\dim\mathfrak{p}=1$ since in this case $\mathrm{so}(\mathfrak{p})=0$ and the three last equations in (19) hold obviously.

**Proposition 4.2.** *If $\dim\mathfrak{p}=1$ then the solutions of Problem 1 are a Kähler Lie algebra $(\mathfrak{h},\varrho,\omega)$, $\phi_{\mathfrak{h}}=0$, $[\ ,\ ]_{\mathfrak{p}}=0$, $\mu=0$ and $\phi_{\mathfrak{p}}(a)\in\mathrm{sp}(\mathfrak{h},\omega)\cap\mathrm{Der}(\mathfrak{h})$ where $a$ is a generator of $\mathfrak{p}$ and $\mathrm{Der}(\mathfrak{h})$ the Lie algebra of derivations of $\mathfrak{h}$.*

Let us solve Problem 1 when $\mathfrak{h}$ is 2-dimensional non abelian.

**Proposition 4.3.** *Let $((\mathfrak{h},\omega,\varrho_{\mathfrak{h}}),(\mathfrak{p},[\ ,\ ]_{\mathfrak{p}},\varrho_{\mathfrak{p}}),\mu,\phi_{\mathfrak{h}},\phi_{\mathfrak{p}})$ be a solution of Problem 1 with $\mathfrak{h}$ is 2-dimensional non abelian. Then there exists an orthonormal basis $\mathbb{B}=(e_1,e_2)$ of $\mathfrak{h}$, $b_0\in\mathfrak{p}$ and two constants $\alpha\neq0$ and $\beta\neq0$ such that:*

(i) $[e_1,e_2]_{\mathfrak{h}}=\alpha e_1$, $\omega=\beta e_1^*\wedge e_2^*$,

(ii) $(\mathfrak{p},[\ ,\ ]_{\mathfrak{p}},\varrho_{\mathfrak{p}})$ *is a Euclidean Lie algebra,*

(iii) $\phi_{\mathfrak{h}}(e_1)=0$, $\phi_{\mathfrak{h}}(e_2)\in\mathrm{Der}(\mathfrak{p})\cap\mathrm{so}(\mathfrak{p})$ *and, for any $a\in\mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a),\mathbb{B})=\begin{pmatrix}0&\varrho_{\mathfrak{p}}(a,b_0)\\0&0\end{pmatrix}$,*

(iv) *for any $a,\ b\in\mathfrak{p}$, $\mu(a,b)=\mu_0(a,b)e_1$ with $\mu_0$ is a 2-cocycle of $(\mathfrak{p},[\ ,\ ]_{\mathfrak{p}})$ satisfying*

$$\mu_0(a,\phi_{\mathfrak{h}}(e_2)b)+\mu_0(\phi_{\mathfrak{h}}(e_2)a,b)=-\varrho_{\mathfrak{p}}([a,b]_{\mathfrak{p}},b_0)-\alpha\mu_0(a,b)\,. \tag{20}$$

**Proof.** Note first that from the third relation in (19) we get that $\phi_{\mathfrak{h}}(\mathfrak{h})$ is a solvable subalgebra of $\mathrm{so}(\mathfrak{p})$ and hence must be abelian. Since $\mathfrak{h}$ is 2-dimensional non abelian then $\dim \phi_{\mathfrak{h}}(\mathfrak{h}) = 1$ and $[\mathfrak{h},\mathfrak{h}] \subset \ker \phi_{\mathfrak{h}}$. So there exists an orthonormal basis $(e_1, e_2)$ of $\mathfrak{h}$ such that $[e_1, e_2]_{\mathfrak{h}} = \alpha e_1$, $\phi_{\mathfrak{h}}(e_1) = 0$ and $\omega = \beta e_1^* \wedge e_2^*$. If we identify the endomorphisms of $\mathfrak{h}$ with their matrices in the basis $(e_1, e_2)$, we get that $\mathrm{sp}(\mathfrak{h}, \omega) = \mathrm{sl}(2, \mathbb{R})$ and there exists $a_0, b_0, c_0 \in \mathfrak{p}$ such that, for any $a \in \mathfrak{p}$,

$$\phi_{\mathfrak{p}}(a) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}.$$

The first equation in (19) is equivalent to

$$\begin{aligned} \alpha\big(\varrho_{\mathfrak{p}}(a_0, a)e_1 + \varrho_{\mathfrak{p}}(c_0, a)e_2\big) = &-\alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 + \alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 \\ &+ \varrho_{\mathfrak{p}}\big(a_0, \phi_{\mathfrak{h}}(e_2)(a)\big)e_1 + \varrho_{\mathfrak{p}}\big(c_0, \phi_{\mathfrak{h}}(e_2)(a)\big)e_2\,, \end{aligned}$$

for any $a \in \mathfrak{p}$. Since $\phi_{\mathfrak{h}}(e_2)$ is sekw-symmetric, this is equivalent to

$$\phi_{\mathfrak{h}}(e_2)(a_0) = -\alpha a_0 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(c_0) = -\alpha c_0\,.$$

This implies that $a_0 = c_0 = 0$. The second equation in (19) implies that $\phi_{\mathfrak{h}}(e_2)$ is a derivation of $[\ ,\ ]_{\mathfrak{p}}$. If we take $u = e_1$ in the forth equation in (19), we get that $[e_1, \mu(a, b)] = 0$, for any $a$, $b \in \mathfrak{p}$ and hence $\mu(a, b) = \mu_0(a, b)e_1$. If we take $u = e_2$ in the forth equation in (19) we get (20). The two last equations are equivalent to $[\ ,\ ]_{\mathfrak{p}}$ is a Lie bracket and $\mu_0$ is 2-cocycle of $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}})$. □

The following proposition gives the solutions of Problem 1 when $\mathfrak{h}$ is 2-dimensional abelian and $\dim \mathfrak{p} = 2$.

**Proposition 4.4.** *Let* $\big((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}}\big)$ *be a solution of Problem 1 with* $\mathfrak{h}$ *is 2-dimensional abelian and* $\dim \mathfrak{p} = 2$. *Then one of the following situations occurs:*

(1) $\phi_{\mathfrak{h}} = 0$, $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ *is a 2-dimensional Euclidean Lie algebra, there exists* $a_0 \in \mathfrak{p}$ *and* $D \in \mathrm{sp}(\mathfrak{h}, \omega)$ *such that, for any* $a \in \mathfrak{p}$, $\phi_{\mathfrak{p}}(a) = \varrho_{\mathfrak{p}}(a_0, a)D$ *and there is no restriction on* $\mu$. *Moreover,* $a_0 \in [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}^{\perp}$ *if* $D \neq 0$.

(2) $\phi_{\mathfrak{h}} = 0$, $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ *is a 2-dimensional non abelian Euclidean Lie algebra,* $\phi_{\mathfrak{p}}$ *identifies* $\mathfrak{p}$ *to a two dimensional subalgebra of* $\mathrm{sp}(\mathfrak{h}, \omega)$ *and there is no restriction on* $\mu$.

(3) $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ *is a Euclidean abelian Lie algebra and there exists an orthonormal basis* $\mathbb{B} = (e_1, e_2)$ *of* $\mathfrak{h}$ *and* $b_0 \in \mathfrak{p}$ *such that* $\omega = \alpha e_1^* \wedge e_2^*$, $\phi_{\mathfrak{h}}(e_1) = 0$, $\phi_{\mathfrak{h}}(e_2) \neq 0$ *and, for any* $a \in \mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$ *and there is no restriction on* $\mu$.

**Proof.** Note first that since $\dim \mathfrak{p} = 2$ the last two equations in (19) hold obviously and $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}})$ is a Lie algebra. We distinguish two cases:

(i) $\phi_{\mathfrak{h}} = 0$. Then (19) is equivalent to $\phi_{\mathfrak{p}}$ is a representation of $\mathfrak{p}$ in $\mathrm{sp}(\mathfrak{h}, \omega) \simeq \mathrm{sl}(2, \mathbb{R})$. Since $\mathrm{sl}(2, \mathbb{R})$ doesn't contain any abelian two dimensional subalgebra, if $\mathfrak{p}$ is an abelian Lie algebra then $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) \leq 1$ and the first situation occurs. If $\mathfrak{p}$ is not abelian then the first or the second situation occurs depending on $\dim \phi_{\mathfrak{p}}(\mathfrak{p})$.

(ii) $\phi_{\mathfrak{h}} \neq 0$. Since $\dim \mathrm{so}(\mathfrak{p}) = 1$ there exists an orthonormal basis $\mathbb{B} = (e_1, e_2)$ of $\mathfrak{h}$ such that $\phi_{\mathfrak{h}}(e_1) = 0$ and $\phi_{\mathfrak{h}}(e_2) \neq 0$. We have $\mathrm{sp}(\mathfrak{h}, \omega) = \mathrm{sl}(2, \mathbb{R})$ and hence, for any $a \in \mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}$. Choose an orthonormal basis $(a_1, a_2)$ of $\mathfrak{p}$. Then there exists $\lambda \neq 0$ such that $\phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2$ and $\phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1$.

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1) = 0\,, \quad a \in \mathfrak{p}\,.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1) = \phi_{\mathfrak{p}}(a_2)(e_1) = 0\,.$$

Then $a_0 = c_0 = 0$ and hence $\phi_{\mathfrak{p}}(a) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$. The second equation in (19) gives

$$\begin{aligned} \phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = {} & [a_1, \phi_{\mathfrak{h}}(e_2)(a_2)]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(e_2)(a_1), a_2]_{\mathfrak{p}} \\ & + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_2)(e_2))(a_2) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_1)(e_2))(a_2)\,, \end{aligned}$$

and hence $\phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = 0$. Thus $[a_1, a_2]_{\mathfrak{p}} = 0$. All the other equations in (19) hold obviously. □

To tackle the last case, we need the determination of 2-dimensional subalgebras of $\mathrm{sl}(2, \mathbb{R})$.

**Proposition 4.5.** *The 2-dimensional subalgebras of* $\mathrm{sl}(2, \mathbb{R})$ *are*

$$\mathfrak{g}_1 = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & -\alpha \end{pmatrix},\ \alpha, \beta \in \mathbb{R} \right\},\quad \mathfrak{g}_2 = \left\{ \begin{pmatrix} \alpha & 0 \\ \beta & -\alpha \end{pmatrix},\ \alpha, \beta \in \mathbb{R} \right\},$$

$$\mathfrak{g}_x = \left\{ \begin{pmatrix} \alpha & \frac{2\beta - \alpha}{x} \\ (\alpha + 2\beta)x & -\alpha \end{pmatrix},\ \alpha, \beta \in \mathbb{R} \right\}$$

*where* $x \in \mathbb{R} \setminus \{0\}$. *Moreover,* $\mathfrak{g}_x = \mathfrak{g}_y$ *if and only if* $x = y$.

**Proof.** Let $\mathfrak{g}$ be a 2-dimensional subalgebra of $\mathrm{sl}(2, \mathbb{R})$. We consider the basis $\mathbb{B} = (h, e, f)$ of $\mathrm{sl}(2, \mathbb{R})$ given by

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},\ f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \text{ and } h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Then

$$[h, e] = 2e\,,\ [h, f] = -2f \quad \text{and} \quad [e, f] = h\,.$$

If $h \in \mathfrak{g}$ then $\mathrm{ad}_h$ leaves $\mathfrak{g}$ invariant. But $\mathrm{ad}_h$ has three eigenvalues $(0, 2, -2)$ with the associated eigenvectors $(h, e, f)$ and hence it restriction to $\mathfrak{g}$ has $(0, 2)$ or $(0, -2)$ as eigenvalues. Thus $\mathfrak{g} = \mathfrak{g}_1$ or $\mathfrak{g} = \mathfrak{g}_2$.

Suppose now that $h \notin \mathfrak{g}$. By using the fact that $\mathrm{sl}(2, \mathbb{R})$ is unimodular, i.e., for any $w \in \mathrm{sl}(2, \mathbb{R})$ $\mathrm{tr}(\mathrm{ad}_w) = 0$, we can choose a basis $(u, v)$ of $\mathfrak{g}$ such that $(u, v, h)$ is a basis of $\mathrm{sl}(2, \mathbb{R})$ and

$$[u, v] = u,\ [h, u] = au + v \quad \text{and} \quad [h, v] = du - av - h\,.$$

If $(x_1, x_2, x_3)$ and $(y_1, y_2, y_3)$ are the coordinates of $u$ and $v$ in $\mathbb{B}$, the brackets above gives

$$\begin{cases} -2(x_1y_3 - x_3y_1) - x_1 = 0\,, \\ 2(x_2y_3 - x_3y_2) - x_2 = 0\,, \\ x_1y_2 - x_2y_1 - x_3 = 0\,, \end{cases} \quad \begin{cases} y_1 = (2-a)x_1\,, \\ y_2 = -(a+2)x_2\,, \\ y_3 = -ax_3\,, \end{cases} \quad \text{and} \quad \begin{cases} dx_1 = (a+2)y_1\,, \\ dx_2 = (a-2)y_2\,, \\ dx_3 = ay_3 + 1\,. \end{cases}$$

Note first that if $x_1 = 0$ then $(x_2, x_3) = (0, 0)$ which impossible so we must have $x_1 \neq 0$ and hence $d = 4 - a^2$. If we replace in the third equation in the second system and the last equation, we get $x_3 = \frac{1}{4}$ and $y_3 = -\frac{a}{4}$. The third equation in the first system gives $x_2 = -\frac{1}{16x_1}$ and hence $y_1 = (2-a)x_1$ and $y_2 = \frac{(a+2)}{16x_1}$. Thus

$$\begin{aligned} \mathfrak{g} &= \operatorname{span}\left\{ \begin{pmatrix} \frac{1}{4} & -\frac{1}{16x_1} \\ x_1 & -\frac{1}{4} \end{pmatrix}, \begin{pmatrix} -\frac{a}{4} & \frac{(a+2)}{16x_1} \\ (2-a)x_1 & \frac{a}{4} \end{pmatrix} \right\} \\ &= \operatorname{span}\left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix} \right\}\,; \quad x = 4x_1\,. \end{aligned}$$

But

$$\begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} = a \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix} + \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix}$$

and hence

$$\mathfrak{g} = \operatorname{span}\left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} \right\} = \mathfrak{g}_x\,.$$

One can check easily that $\mathfrak{g}_x = \mathfrak{g}_y$ if and only if $x = y$. This completes the proof. □

The following two propositions give the solutions of Problem 1 when $\mathfrak{h}$ is 2-dimensional abelian and $\dim \mathfrak{p} = 3$.

**Proposition 4.6.** *Let $((\mathfrak{h}, \omega, \varrho_\mathfrak{h}), (\mathfrak{p}, [\ ,\ ]_\mathfrak{p}, \varrho_\mathfrak{p}), \mu, \phi_\mathfrak{h}, \phi_\mathfrak{p})$ be a solution of Problem 1 with $\mathfrak{h}$ is 2-dimensional abelian and $\dim \mathfrak{p} = 3$ and $\phi_\mathfrak{h} = 0$. Then one of the following situations occurs:*

(i) *$(\mathfrak{p}, [\ ,\ ]_\mathfrak{p}, \varrho_\mathfrak{p})$ is 3-dimensional Euclidean Lie algebra, $\phi_\mathfrak{p} = 0$ and $\mu$ is 2-cocycle for the trivial representation.*

(ii) *$\phi_\mathfrak{p}$ is an isomorphism of Lie algebras between $(\mathfrak{p}, [\ ,\ ]_\mathfrak{p})$ and $\mathrm{sl}(2, \mathbb{R})$ and there exists an endomorphism $L\colon \mathfrak{p} \longrightarrow \mathfrak{h}$ such that for any $a$, $b \in \mathfrak{p}$,*

$$\mu(a, b) = \phi_\mathfrak{p}(a)(L(b)) - \phi_\mathfrak{p}(b)(L(a)) - L([a, b]_\mathfrak{p})\,.$$

(iii) *There exists a basis $\mathbb{B}_\mathfrak{p} = (a_1, a_2, a_3)$ of $\mathfrak{p}$, $\alpha \neq 0$, $\beta \neq 0$, $\gamma$, $\tau \in \mathbb{R}$ such that $[\ ,\ ]_\mathfrak{p}$ has one of the two following forms*

$$\begin{cases} [a_1, a_2]_\mathfrak{p} = 0\,,\ [a_1, a_3]_\mathfrak{p} = \beta a_1\,, \\ [a_2, a_3]_\mathfrak{p} = \gamma a_1 + \alpha a_2\,,\ \alpha \neq 0,\ \beta \neq 0 \\ M(\varrho_\mathfrak{p}, \mathbb{B}_\mathfrak{p}) = \mathrm{I}_3 \end{cases} \quad \text{or} \quad \begin{cases} [a_1, a_2]_\mathfrak{p} = [a_1, a_3]_\mathfrak{p} = 0\,, \\ [a_2, a_3]_\mathfrak{p} = \alpha a_2\,,\ \alpha \neq 0\,, \\ M(\varrho_\mathfrak{p}, \mathbb{B}_\mathfrak{p}) = \begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}\,. \end{cases}$$

*In both cases, there exists an orthonormal basis* $\mathbb{B}_{\mathfrak{h}} = (e_1, e_2)$ *of* $\mathfrak{h}$, $x \neq 0$, $u \neq 0$ *and* $v \in \mathbb{R}$ *such that* $\phi_{\mathfrak{p}}$ *has one of the following forms*

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \\ \mathcal{M}(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0\,, \end{cases} \qquad \begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0\,, \end{cases} \quad or$$

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} v & -\frac{2v+\alpha}{2x} \\ \frac{2v-\alpha}{2}x & -v \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0\,. \end{cases}$$

*Moreover,* $\mu$ *is a 2-cocycle for* $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

(iv) *There exists an orthonormal basis* $\mathbb{B} = (a_1, a_2, a_3)$ *of* $\mathfrak{p}$ *such that* $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$, $\phi_{\mathfrak{p}}(a_3)$ *is a non zero element of* $\mathrm{sp}(\mathfrak{h}, \omega)$ *and*

$$\begin{cases} [a_1, a_2]_{\mathfrak{p}} = 0\,,\ [a_1, a_3]_{\mathfrak{p}} = \beta a_1 + \rho a_2\,, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2\,, \end{cases} \quad or \quad \begin{cases} [a_1, a_2]_{\mathfrak{p}} = \alpha a_2\,,\ [a_1, a_3]_{\mathfrak{p}} = \rho a_2\,, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_2, \alpha \neq 0\,. \end{cases}$$

*Moreover,* $\mu$ *is a 2-cocycle for* $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

**Proof.** In this case, (19) is equivalent to $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}})$ is a Lie algebra and $\phi_{\mathfrak{p}}$ is a representation and $\mu$ is a 2-cocycle of $(\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

We distinguish four cases:

(1) $\phi_{\mathfrak{p}} = 0$ and the case $(i)$ occurs.

(2) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 3$ and hence $\mathfrak{p}$ is isomorphic to $\mathrm{sp}(\mathfrak{h}, \omega) \simeq \mathrm{sl}(2, \mathbb{R})$ and hence $\mu$ is a coboundary. Thus $(ii)$ occurs.

(3) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 2$ then $\ker \phi_{\mathfrak{p}}$ is a one dimensional ideal of $\mathfrak{p}$. But $\phi_{\mathfrak{p}}(\mathfrak{p})$ is a 2-dimensional subalgebra of $\mathrm{sp}(\mathfrak{h}, \omega) \simeq \mathrm{sl}(2, \mathbb{R})$, therefore it is non abelian so $\mathfrak{p}/\ker \mathfrak{p}$ is non abelian.

If $\ker \mathfrak{p} \subset [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$ then $\dim [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = 2$ so there exists an orthonormal basis $(a_1, a_2, a_3)$ of $\mathfrak{p}$ such that $a_1 \in \ker \mathfrak{p}$ and

$$[a_1, a_2]_{\mathfrak{p}} = \xi a_1,\ [a_1, a_3]_{\mathfrak{p}} = \beta a_1 \quad \text{and} \quad [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2,\ \alpha \neq 0, \beta \neq 0$$

and we must have $\xi = 0$ in order to have the Jacobi identity.

If $\ker \mathfrak{p} \not\subset [\mathfrak{p}, \mathfrak{p}]$ then $\ker \mathfrak{p} \subset Z(\mathfrak{p})$ and $\dim [\mathfrak{p}, \mathfrak{p}] = 1$. Then there exits a basis $(a_1, a_2, a_3)$ of $\mathfrak{p}$ such that $a_1 \in \ker \mathfrak{p}$, $a_2 \in [\mathfrak{p}, \mathfrak{p}]$, $a_3 \in \{a_1, a_2\}^{\perp}$ and

$$[a_2, a_3]_{\mathfrak{p}} = \alpha a_2,\ [a_3, a_1]_{\mathfrak{p}} = [a_1, a_2]_{\mathfrak{p}} = 0,\ \alpha \neq 0.$$

The matrix of $\varrho_{\mathfrak{p}}$ in $(a_1, a_2, a_3)$ is given by

$$\begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We choose an orthonormal basis $(e_1, e_2)$ of $\mathfrak{h}$ and identify $\mathrm{sp}(\mathfrak{h}, \omega)$ to $\mathrm{sl}(2, \mathbb{R})$. Now $\phi_{\mathfrak{p}}(\mathfrak{p}) = \{\phi_p(a_2), \phi_p(a_3)\}$ is a subalgebra of $\mathrm{sl}(2, \mathbb{R})$ and, according to Proposition 4.5, $\phi_{\mathfrak{p}}(\mathfrak{p}) = \mathfrak{g}_1$, $\mathfrak{g}_2$ or $\mathfrak{g}_x$. But

$$[\mathfrak{g}_1, \mathfrak{g}_1] = \mathbb{R}e, [\mathfrak{g}_2, \mathfrak{g}_2] = \mathbb{R}f \quad \text{and} \quad [\mathfrak{g}_x, \mathfrak{g}_x] = \left\{ \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix} \right\}.$$

So in order for $\phi_{\mathfrak{p}}$ to be a representation we must have

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \quad \text{and} \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \ \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

or

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \ \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} p & -\frac{2p+\alpha}{2x} \\ \frac{2p-\alpha}{2}x & -p \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0.$$

(4) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 1$ then $\ker \phi_{\mathfrak{p}}$ is a two dimensional ideal of $\mathfrak{p}$. Then there exists an orthonormal basis $(a_1, a_2, a_3)$ of $\mathfrak{p}$ such that

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_2, \ [a_3, a_1]_{\mathfrak{p}} = pa_1 + qa_2 \quad \text{and} \quad [a_3, a_2]_{\mathfrak{p}} = ra_1 + sa_2.$$

The Jacobi identity gives $\alpha = 0$ or $(p, r) = (0, 0)$. We take $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$ and $\phi_{\mathfrak{p}}(a_3) \in \mathrm{sl}(2, \mathbb{R})$. □

**Proposition 4.7.** *Let* $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\ ,\ ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ *be a solution of Problem 1 with* $\mathfrak{h}$ *is 2-dimensional abelian,* $\dim \mathfrak{p} = 3$ *and* $\phi_{\mathfrak{h}} \neq 0$. *Then there exists an orthonormal basis* $(e_1, e_2)$ *of* $\mathfrak{h}$, *an orthonormal basis* $(a_1, a_2, a_3)$ *of* $\mathfrak{p}$, $\lambda > 0$, $\alpha, p, q, \mu_1, \mu_2, \mu_3 \in \mathbb{R}$ *such that*

$$\phi_{\mathfrak{h}}(e_1) = 0, \ \phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2, \ \phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1 \quad \textit{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3) = 0,$$

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_3, \ [a_1, a_3]_{\mathfrak{p}} = pa_1 + qa_2, [a_2, a_3]_{\mathfrak{p}} = -qa_1 + pa_2 \quad \textit{and}$$

$$\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}, \quad i = 1, 2, 3$$

*and one of the following situations occurs:*

(i) $p \neq 0$, $\alpha = 0$ *and*

$$\mu(a_1, a_2) = 0, \ \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \ \textit{ and } \ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(ii) $p = 0$, $\mu_3 \neq 0$, $\alpha = 0$ *and*

$$\mu(a_1, a_2) = ce_1, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \ \textit{ and } \ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(iii) $p=0$, $\mu_3=0$ *and*

$$\mu(a_1,a_2)=c_1e_1+c_2e_2, \mu(a_2,a_3)=-\lambda^{-1}(p\mu_1+q\mu_2)e_1 \quad \textit{and}$$
$$\mu(a_1,a_3)=\lambda^{-1}(-q\mu_1+p\mu_2)e_1\,.$$

**Proof.** Since $\phi_{\mathfrak{h}}\neq 0$ then $\phi_{\mathfrak{h}}(\mathfrak{h})$ is a non trivial abelian subalgebra of $\mathrm{so}(\mathfrak{p})$ and hence it must be one dimensional. Then there exists an orthonormal basis $(e_1,e_2)$ of $\mathfrak{h}$ and an orthonormal basis $(a_1,a_2,a_3)$ of $\mathfrak{p}$ and $\lambda>0$ such that $\phi_{\mathfrak{h}}(e_1)=0$ and

$$\phi_{\mathfrak{h}}(e_2)(a_1)=\lambda a_2,\ \phi_{\mathfrak{h}}(e_2)(a_2)=-\lambda a_1 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3)=0\,.$$

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1)=0\,, \quad a\in\mathfrak{p}\,.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1)=\phi_{\mathfrak{p}}(a_2)(e_1)=0\,.$$

Thus $\phi_{\mathfrak{p}}(a_i)=\begin{pmatrix}0 & \mu_i\\ 0 & 0\end{pmatrix}$ for $i=1,2$ and $\phi_{\mathfrak{p}}(a_3)=\begin{pmatrix}u & v\\ w & -u\end{pmatrix}$. Consider now the second equation in (19)

$$\phi_{\mathfrak{h}}(u)([a,b]_{\mathfrak{p}})=[a,\phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}}+[\phi_{\mathfrak{h}}(u)(a),b]_{\mathfrak{p}}+\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a)-\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b)\,.$$

This equation is obviously true when $u=e_1$ and $(a,b)=(a_1,a_2)$. For $u=e_1$ and $(a,b)=(a_1,a_3)$, we get

$$\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_3)(e_1))(a_1)=0$$

and hence $w=0$.

For $u=e_2$ and $(a,b)=(a_1,a_2)$, we get $\phi_{\mathfrak{h}}(e_2)([a_1,a_2]_{\mathfrak{p}})=0$ and hence $[a_1,a_2]_{\mathfrak{p}}=\alpha a_3$.

For $u=e_2$ and $(a,b)=(a_1,a_3)$ or $(a,b)=(a_2,a_3)$ , we get

$$\phi_{\mathfrak{h}}(e_2)([a_1,a_3]_{\mathfrak{p}})=\lambda[a_2,a_3]_{\mathfrak{p}}-\lambda u a_2 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)([a_2,a_3]_{\mathfrak{p}})=-\lambda[a_1,a_3]_{\mathfrak{p}}+\lambda u a_1.$$

This implies that $[a_1,a_3]_{\mathfrak{p}},[a_2,a_3]_{\mathfrak{p}}\in\mathrm{span}\{a_1,a_2\}$ and hence

$$[a_1,a_3]_{\mathfrak{p}}=pa_1+qa_2 \quad \text{and} \quad [a_2,a_3]_{\mathfrak{p}}=ra_1+sa_2.$$

So

$$\begin{cases}\lambda(pa_2-qa_1)=\lambda(ra_1+sa_2-ua_2)\,,\\ \lambda(ra_2-sa_1)=-\lambda(pa_1+qa_2-ua_1)\,.\end{cases}$$

This is equivalent to

$$u=0, p=s \quad \text{and} \quad r=-q\,.$$

To summarize, we get

$$[a_1,a_2]_{\mathfrak{p}}=\alpha a_3,\ [a_1,a_3]_{\mathfrak{p}}=pa_1+qa_2, [a_2,a_3]_{\mathfrak{p}}=-qa_1+pa_2 \quad \text{and}$$
$$\phi_{\mathfrak{p}}(a_i)=\begin{pmatrix}0 & \mu_i\\ 0 & 0\end{pmatrix}\,.$$

Let consider now the fourth equation in (19)

$$\phi_{\mathfrak{p}}([a,b]_{\mathfrak{p}})(u)=[\phi_{\mathfrak{p}}(a),\phi_{\mathfrak{p}}(b)](u)+[u,\mu(a,b)]_{\mathfrak{h}}-\mu(a,\phi_{\mathfrak{h}}(u)(b))-\mu(\phi_{\mathfrak{h}}(u)(a),b).$$

This equation is obviously true for $u=e_1$.

For $u = e_2$ and $(a,b) = (a_1,a_2)$, $(a,b) = (a_1,a_3)$ or $(a,b) = (a_2,a_3)$, we get

$$\begin{cases} \alpha\mu_3 = 0, \\ (p\mu_1 + q\mu_2)e_1 = -\lambda\mu(a_2,a_3)\,, \\ (-q\mu_1 + p\mu_2)e_1 = \lambda\mu(a_1,a_3)\,. \end{cases}$$

The last two equations are equivalent to

$$\phi_{\mathfrak{p}}(a_3)(\mu(a_1,a_2)) = -2p\mu(a_1,a_2) \quad \text{and} \quad p[a_1,a_2]_{\mathfrak{p}} = 0\,.$$

• $p \neq 0$ then

$$\alpha = 0, \mu(a_1,a_2) = 0,\ \mu(a_2,a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and}$$
$$\mu(a_1,a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1\,.$$

• $p = 0$ and $\mu_3 \neq 0$ then $\alpha = 0$ and

$$\mu(a_1,a_2) = ce_1, \mu(a_2,a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and}$$
$$\mu(a_1,a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1\,.$$

• $p = 0$ and $\mu_3 = 0$ then

$$\mu(a_1,a_2) = c_1e_1 + c_2e_2, \mu(a_2,a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and}$$
$$\mu(a_1,a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1\,.$$

□

By using Propositions 4.2–4.7, we can give all the Riemann-Poisson Lie algebras of dimension 3, 4 or 5.

Let $(\mathfrak{g}, [\ ,\ ], \varrho, r)$ be a Riemann-Poisson Lie algebra of dimension less or equal to 5. According to what above then $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ and the Lie bracket on $\mathfrak{g}$ is given by (18) and $((\mathfrak{h},\omega,\varrho_{\mathfrak{h}}),(\mathfrak{p},[\ ,\ ]_{\mathfrak{p}},\varrho_{\mathfrak{p}}),\mu,\phi_{\mathfrak{h}},\phi_{\mathfrak{p}})$ are solutions of Problem 1.

• $\dim\mathfrak{g} = 3$. In this case $\dim\mathfrak{h} = 2$ and $\dim\mathfrak{p} = 1$ and, by applying Proposition 4.2, the Lie bracket of $\mathfrak{g}$, $\varrho$ and $r$ are given in Table 1, where $e^{12} = e_1 \wedge e_2$.

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_1,e_2] = ae_1, [e_3,e_2] = be_1$ | $\alpha e^{12}$ | $\mathrm{I}_3$ | $a \neq 0, \alpha \neq 0$ |
| $[e_3,e_1] = -be_1 + ce_2, [e_3,e_2] = de_1 + be_2$ | $\alpha e^{12}$ | $\mathrm{I}_3$ | $\alpha \neq 0$ |

TAB. 1. Three dimensional Riemann-Poisson Lie algebras

• $\dim\mathfrak{g} = 4$. We have three cases:

($c41$) $\dim\mathfrak{h} = 2$, $\dim\mathfrak{p} = 2$ and $\mathfrak{h}$ is non abelian and we can apply Proposition 4.3 to get the Lie brackets on $\mathfrak{g}$, $\varrho$ and $r$. They are described in rows 1 and 2 in Table 2.

($c42$) $\dim\mathfrak{h} = 2$, $\dim\mathfrak{p} = 2$ and $\mathfrak{h}$ is abelian and we can apply Propositions 4.4 and 4.5 to get the Lie brackets on $\mathfrak{g}$, $\varrho$ and $r$. They are described in rows 3 and 8 in Table 2.

($c43$) $\dim\mathfrak{h} = 4$. In this case $\mathfrak{g}$ is a Kähler Lie algebra. We have used [12] to derive all four dimensional Kähler Lie algebra together with their symplectic derivations. The results are given in Table 3. The notation $\mathrm{Der}^s(\mathfrak{h})$ stands

for the vector spaces of derivations which are skew-symmetric with respect the symplectic form. The vector space $\mathrm{Der}^s(\mathfrak{h})$ is described by a family of generators and $E_{ij}$ is the matrix with 1 in the $i$ row and $j$ column and 0 elsewhere.

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_1,e_2]=ae_1,\ [e_3,e_2]=be_1+ce_4,$ $[e_4,e_2]=de_1-ce_3$ | $\alpha e^{12}$ | $\mathrm{I}_4$ | $a\neq 0,\alpha\neq 0$ |
| $[e_1,e_2]=ae_1,\ [e_3,e_2]=be_1,$ $[e_4,e_2]=de_1,[e_3,e_4]=ce_3-a^{-1}cbe_1$ | $\alpha e^{12}$ | $\mathrm{I}_4$ | $\alpha ac\neq 0,$ |
| $[e_3,e_4]=ae_1+be_2$ | $\alpha e^{12}$ | $\mathrm{I}_4$ | $\alpha\neq 0$ |
| $[e_3,e_4]=ae_1+be_2+ce_3,\ [e_4,e_1]=xe_1+ye_2,$ $[e_4,e_2]=ze_1-xe_2$ | $\alpha e^{12}$ | $\mathrm{I}_4$ | $\alpha\neq 0$ |
| $[e_3,e_4]=ae_1+be_2+2e_4,\ [e_3,e_1]=e_1,$ $[e_3,e_2]=-e_2,[e_4,e_2]=e_1$ | $\alpha e^{12}$ | $\mathrm{Diag}\left(1,1,\begin{pmatrix}\mu&\nu\\ \nu&\rho\end{pmatrix}\right)$ | $\alpha\neq 0,\mu,\rho>0$ $\mu\rho>\nu^2$ |
| $[e_3,e_4]=ae_1+be_2-2e_4,\ [e_3,e_1]=e_1,$ $[e_3,e_2]=-e_2,[e_4,e_1]=e_2$ | $\alpha e^{12}$ | $\mathrm{Diag}\left(1,1,\begin{pmatrix}\mu&\nu\\ \nu&\rho\end{pmatrix}\right)$ | $\alpha\neq 0,\mu,\rho>0$ $\mu\rho>\nu^2$ |
| $[e_3,e_4]=ae_1+be_2-2e_3,\ [e_3,e_1]=e_1+xe_2,$ $[e_3,e_2]=-\frac{1}{x}e_1-e_2,[e_4,e_1]=xe_2,[e_4,e_2]=\frac{1}{x}e_1$ | $\alpha e^{12}$ | $\mathrm{Diag}\left(1,1,\begin{pmatrix}\mu&\nu\\ \nu&\rho\end{pmatrix}\right)$ | $\alpha\neq 0,\mu,\rho>0$ $\mu\rho>\nu^2,x\neq 0$ |
| $[e_3,e_4]=ae_1+be_2,\ [e_3,e_2]=xe_1+ye_4,$ $[e_4,e_2]=ze_1-ye_3$ | $\alpha e^{12}$ | $\mathrm{I}_4$ | $\alpha y\neq 0$ |

TAB. 2. Four dimensional Riemann-Poisson Lie algebras of rank 2

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | $\mathrm{Der}^s(\mathfrak{h})$ |
|---|---|---|---|
| $[e_1,e_2]=e_2,$ | $\alpha e^{12}+\beta e^{34}$ | $\mathrm{Diag}(a,b,c,d)$ | $\{E_{21},E_{33}-E_{44},E_{43},E_{34}\}$ |
| $[e_1,e_2]=-e_3,[e_1,e_3]=e_2,$ | $\alpha e^{14}+\beta e^{23}$ | $\mathrm{Diag}(a,b,b,c)$ | $\{E_{23}-E_{32},E_{41}\}$ |
| $[e_1,e_2]=e_2,[e_3,e_4]=e_4,$ | $\alpha e^{12}+\beta e^{34}$ | $\mathrm{Diag}(a,b,c,d)$ | $\{E_{21},E_{43}\}$ |
| $[e_4,e_1]=e_1,[e_4,e_2]=-\delta e_3,$ $[e_4,e_3]=\delta e_2$ | $\alpha e^{14}+\beta e^{23}$ | $\mathrm{Diag}(a,b,b,c)$ | $\{E_{14},E_{23}-E_{32}\}$ |
| $[e_1,e_2]=e_3,[e_4,e_3]=e_3,$ $[e_4,e_1]=\frac{1}{2}e_1,[e_4,e_2]=\frac{1}{2}e_2,,$ | $\alpha(e^{12}-e^{34})$ | $\mathrm{Diag}(a,\mu b,\mu a,b)$ | $\{E_{34},E_{22}-E_{11},E_{12}+E_{21}\}$ |
| $[e_1,e_2]=e_3,[e_4,e_3]=e_3,$ $[e_4,e_1]=2e_1,[e_4,e_2]=-e_2,$ | $\alpha(e^{23}+e^{14})$ | $\mathrm{Diag}(a,a,2a,2a)$ | $\{2E_{14}-E_{32}\}$ |
| $[e_1,e_2]=e_3,[e_4,e_3]=e_3,$ $[e_4,e_1]=\frac{1}{2}e_1-e_2,$ $[e_4,e_2]=e_1+\frac{1}{2}e_2,$ | $\alpha(e^{12}-e^{34})$ | $\mathrm{Diag}(a,a,a,a)$ | $\{E_{34},E_{12}-E_{21}\}$ |

TAB. 3. Four-dimensional Kähler Lie algebras and their symplectic derivations, $a,b,c,d>0$,$\alpha\beta\neq 0$

- $\dim\mathfrak{g}=5$. We have:

(c51) $\dim\mathfrak{h}=4$ and $\mathfrak{h}$ abelian and hence a symplectic vector space. We can apply Proposition 4.2 and $\mathfrak{g}$ is semi-direct product.

(c52) $\dim\mathfrak{h}=4$ and $\mathfrak{h}$ non abelian. We can apply Proposition 4.2 and Table 3 to get the Lie brackets on $\mathfrak{g}$, $\varrho$ and $r$. The result is summarized in Table 4.

(c53) $\dim\mathfrak{h}=2$ and $\mathfrak{h}$ non abelian. We apply Proposition 4.3. In this case $(\mathfrak{p},[\ ,\ ]_{\mathfrak{p}},\varrho_{\mathfrak{p}})$ is a 3-dimensional Euclidean Lie algebra and one must compute $\mathrm{Der}(\mathfrak{p})\cap\mathrm{so}(\mathfrak{p})$ and solve (20). Three dimensional Euclidean Lie algebras were classified in [9]. For each of them we have computed $\mathrm{Der}(\mathfrak{p})\cap\mathrm{so}(\mathfrak{p})$

and solved (20) by using Maple. The result is summarized in Table 5 when $\mathfrak{p}$ is unimodular and Table 6 when $\mathfrak{p}$ is nonunimodular.

($c54$) $\dim \mathfrak{h} = 2$ and $\mathfrak{h}$ abelian and $\phi_{\mathfrak{h}} = 0$. We apply Proposition 4.6 and we perform all the needed computations. We use the classification of 3-dimensional Euclidean Lie algebras given in [9]. The results are given in Tables 7-8.

($c55$) $\dim \mathfrak{h} = 2$ and $\mathfrak{h}$ abelian and $\phi_{\mathfrak{h}} \neq 0$. We apply Proposition 4.7 and we perform all the needed computations. The results are given in Table 9.

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_1,e_2]=e_2, [e_5,e_1]=xe_2,$<br>$[e_5,e_3]=ye_3+te_4, [e_5,e_4]=ze_3-ye_4$ | $\alpha e^{12}+\beta e^{34}$ | $\mathrm{Diag}(a,b,c,d,e)$ | $\alpha\beta\neq 0$<br>$a,b,c,d,e>0$ |
| $[e_1,e_2]=-e_3, [e_1,e_3]=e_2,$<br>$[e_5,e_1]=ye_4, [e_5,e_2]=-xe_3, [e_5,e_3]=xe_2$ | $\alpha e^{14}+\beta e^{23}$ | $\mathrm{Diag}(a,b,b,c,d)$ | $\alpha\beta\neq 0$<br>$a,b,c,d>0$ |
| $[e_1,e_2]=e_2, [e_3,e_4]=e_4,$<br>$[e_5,e_1]=xe_2, [e_5,e_3]=ye_4$ | $\alpha e^{12}+\beta e^{34}$ | $\mathrm{Diag}(a,b,c,d,e)$ | $\alpha\beta\neq 0$<br>$a,b,c,d,e>0$ |
| $[e_4,e_1]=e_1, [e_4,e_2]=-\delta e_3, [e_4,e_3]=\delta e_2$<br>$[e_5,e_2]=-ye_3, [e_5,e_3]=ye_2, [e_5,e_4]=xe_1$ | $\alpha e^{14}+\beta e^{23}$ | $\mathrm{Diag}(a,b,b,c,d)$ | $\alpha\beta\neq 0, \delta>0$<br>$a,b,c,d>0$ |
| $[e_1,e_2]=e_3, [e_4,e_3]=e_3, [e_4,e_1]=\frac{1}{2}e_1$<br>$[e_4,e_2]=\frac{1}{2}e_2, [e_5,e_1]=xe_1+ye_2,$<br>$[e_5,e_2]=ye_1-xe_2, [e_5,e_4]=ze_3$ | $\alpha(e^{12}-e^{34})$ | $\mathrm{Diag}(a,\mu b,\mu a,b,c)$ | $\alpha\neq 0$<br>$a,b,c,\mu>0$ |
| $[e_1,e_2]=e_3, [e_4,e_3]=e_3, [e_4,e_1]=2e_1$<br>$[e_4,e_2]=-e_2, [e_5,e_2]=xe_3, [e_5,e_4]=-2xe_1$ | $\alpha(e^{23}+e^{14})$ | $\mathrm{Diag}(a,a,2a,2a,b)$ | $\alpha\neq 0$<br>$a,b>0$ |
| $[e_1,e_2]=e_3, [e_4,e_3]=e_3, [e_4,e_1]=\frac{1}{2}e_1-e_2$<br>$[e_4,e_2]=e_1+\frac{1}{2}e_2, [e_5,e_1]=-xe_2, [e_5,e_2]=xe_1$<br>$[e_5,e_4]=ye_3$ | $\alpha(e^{12}-e^{34})$ | $\mathrm{Diag}(a,a,a,a,b)$ | $\alpha\neq 0$<br>$a,b>0$ |

TAB. 4. Five-dimensional Riemann-Poisson Lie algebras of rank 4

| Non vanishing Lie brackets | $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1-ce_4, [e_4,e_2]=d\mu e_1+ce_3$<br>$[e_5,e_2]=fe_1, [e_3,e_4]=-fe_1+e_5$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\mu,\mu,1)$ | $c\alpha\neq 0$<br>$\mu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=be_1, [e_4,e_2]=ce_1$<br>$[e_5,e_2]=d\mu e_1, [e_3,e_5]=be_1-e_3, [e_4,e_5]=-ce_1+e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,1,1,\mu)$ | $\alpha\neq 0$<br>$\mu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=(b+c)e_1, [e_4,e_2]=(cx+b)e_1$<br>$[e_5,e_2]=d\mu e_1, [e_3,e_5]=(b+c)e_1-e_3,$<br>$[e_4,e_5]=-(xc+b)e_1+e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\begin{pmatrix}1&1\\1&x\end{pmatrix},\mu)$ | $\alpha\neq 0$<br>$\mu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=be_1, [e_4,e_2]=c\mu e_1$<br>$[e_5,e_2]=d\nu e_1, [e_3,e_5]=-\mu ce_1+e_4, [e_4,e_5]=be_1-e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,1,\mu,\nu)$ | $\alpha\neq 0$<br>$\mu,\nu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1, [e_4,e_2]=c\nu e_1$<br>$[e_5,e_2]=d\rho e_1, [e_3,e_4]=-2\rho de_1+2e_5,$<br>$[e_3,e_5]=2\nu ce_1-2e_4, [e_4,e_5]=2\mu be_1-2e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\xi,\mu,\nu,\rho)$ | $\alpha\neq 0, \nu\neq\rho$<br>$\mu,\nu,\rho,\xi>0$<br>$\mu\neq\nu, \mu\neq\rho$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1, [e_4,e_2]=c\nu e_1-\lambda e_5$<br>$[e_5,e_2]=d\nu e_1+\lambda e_4, [e_3,e_4]=-\frac{2\nu(\lambda c+d)}{1+\lambda^2}e_1+2e_5,$<br>$[e_3,e_5]=\frac{2\nu(c-\lambda d)}{1+\lambda^2}e_1-2e_4, [e_4,e_5]=2\mu be_1-2e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\mu,\nu,\nu)$ | $\lambda\alpha\neq 0$<br>$\mu,\nu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1, [e_4,e_2]=c\nu e_1$<br>$[e_5,e_2]=d\rho e_1, [e_3,e_4]=-\rho de_1+e_5,$<br>$[e_3,e_5]=\nu ce_1-e_4, [e_4,e_5]=-\mu be_1+e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\xi,\mu,\nu,\rho)$ | $\alpha\neq 0, \nu\neq\rho$<br>$\mu,\nu,\rho,\xi>0$<br>$\mu\neq\nu, \mu\neq\rho$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1, [e_4,e_2]=c\nu e_1-\lambda e_5$<br>$[e_5,e_2]=d\nu e_1+\lambda e_4, [e_3,e_4]=-\frac{\nu(\lambda c+d)}{1+\lambda^2}e_1+e_5,$<br>$[e_3,e_5]=\frac{\nu(c-\lambda d)}{1+\lambda^2}e_1-e_4, [e_4,e_5]=-\mu be_1+e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\mu,\nu,\nu)$ | $\lambda\alpha\neq 0$<br>$\mu,\nu,\rho>0$ |
| $[e_1,e_2]=e_1, [e_3,e_2]=b\mu e_1-ue_4-ve_5,$<br>$[e_4,e_2]=c\mu e_1+ue_3-we_5, [e_5,e_2]=d\mu e_1+ve_3+we_4,$<br>$[e_3,e_4]=xe_1+e_5, [e_3,e_5]=ye_1-e_4, [e_4,e_5]=ze_1+e_3$<br>$x=-\frac{\mu(buw-cuv+du^2+bv+cw+d)}{1+u^2+v^2+w^2}$<br>$y=\frac{\mu(-bvw+cv^2-duw+bu-dw+c)}{1+u^2+v^2+w^2}$<br>$z=-\frac{\mu(bw^2-cvw+duw-cu-dv+b)}{1+u^2+v^2+w^2}$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\mu,\mu,\mu)$ | $\alpha\neq 0$<br>$\mu,\rho>0$ |

TAB. 5. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and unimodular complement

| Non vanishing Lie brackets | $r$ | Matrix of $\varrho$ |
|---|---|---|
| $[e_1,e_2]=e_1,[e_3,e_2]=(f+c\lambda+f\lambda^2)e_1-\lambda e_4,\lambda\neq 0,$<br>$[e_4,e_2]=ce_1+\lambda e_3,[e_5,e_2]=d\mu e_1,$<br>$[e_3,e_5]=fe_1-e_3,[e_4,e_5]=(\lambda f+c)e_1-e_4,$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,1,1,\mu)$<br>$\mu,\rho>0$ |
| $[e_1,e_2]=e_1,[e_3,e_2]=be_1,[e_4,e_2]=c\mu e_1,$<br>$[e_5,e_2]=d\nu e_1,[e_3,e_5]=\mu ce_1-e_4,$<br>$[e_4,e_5]=(-fb+2\mu c)e_1+fe_3-2e_4,f=1$ or $f\leq 0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,1,\mu,\nu)$<br>$0<\mu<\lvert f\rvert,\rho>0$ |
| $[e_1,e_2]=e_1,[e_3,e_2]=(b+c\mu)e_1,[e_4,e_2]=(c+b\mu)e_1,$<br>$[e_5,e_2]=d\nu e_1,[e_3,e_5]=(\mu b+c)e_1-e_4,$<br>$[e_4,e_5]=((2-\mu)c+(2\mu-1)b)e_1+e_3-2e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\begin{pmatrix}1&\mu\\ \mu&1\end{pmatrix},\nu)$<br>$\mu,\nu,\rho>0$ |
| $[e_1,e_2]=e_1,[e_3,e_2]=(b+c)e_1,[e_4,e_2]=(b+c\mu)e_1,$<br>$[e_5,e_2]=d\nu e_1,[e_3,e_5]=(b+c\mu)e_1-e_4,$<br>$[e_4,e_5]=((2-f)b+(2\mu-f)c)e_1+fe_3-2e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\begin{pmatrix}1&1\\ 1&\mu\end{pmatrix},\nu)$<br>$\nu,\rho>0,c>\mu>1$ |
| $[e_1,e_2]=e_1,[e_3,e_2]=(b+\frac{1}{2}c)e_1,[e_4,e_2]=(c+\frac{1}{2}b)e_1,$<br>$[e_5,e_2]=d\nu e_1,[e_3,e_5]=(c+\frac{1}{2}b)e_1-e_4,$<br>$[e_4,e_5]=(b+2c)e_1-2e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,\rho,\begin{pmatrix}1&\frac{1}{2}\\ \frac{1}{2}&1\end{pmatrix},\nu)$<br>$\rho,\nu>0$ |
| $[e_1,e_2]=e_1,[e_3,e_2]=xe_1,[e_4,e_2]=ye_1,$<br>$[e_5,e_2]=d\nu e_1,[e_3,e_5]=ze_1-e_4,$<br>$[e_4,e_5]=te_1+fe_3-2e_4,0<f<1,$<br>$x=\frac{((\mu+1)b+(\mu-1)c)f-2b}{2f^2(f-1)},y=z=\frac{(\mu-1)(cf+b)}{2f(f-1)}$<br>$t=\frac{(1-\mu)cf+((f-2)\mu+f)b}{2f(1-f)}$ | $\alpha e^{12}$ | $A^tBA$<br>$A=\begin{pmatrix}\frac{1+s}{-2fs}&-\frac{1}{2s}&0\\ \frac{1-s}{2fs}&\frac{1}{2s}&\\ 0&0&1\end{pmatrix}$<br>$B=\mathrm{Diag}(1,\rho,\begin{pmatrix}1&\mu\\ \mu&1\end{pmatrix},\nu)$<br>$\nu,\rho>0$<br>$s=\sqrt{1-f},\ 0\leq\mu<1$ |

TAB. 6. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and non unimodular complement. ($\alpha\neq 0$)

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_3,e_4]=ae_1+be_2+e_5,[e_3,e_5]=ce_1+de_2$<br>$[e_4,e_5]=fe_1+ge_2$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\mu,\mu,1)$ | $\alpha\neq 0$<br>$\mu>0$ |
| $[e_3,e_4]=ae_1+be_2,[e_3,e_5]=ce_1+de_2-e_3$<br>$[e_4,e_5]=fe_1+ge_2+e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,\mu)$<br>$\mathrm{Diag}(1,1,\begin{pmatrix}1&1\\ 1&x\end{pmatrix},\mu)$ | $\alpha\neq 0$<br>$\mu>0$ |
| $[e_3,e_4]=ae_1+be_2,[e_3,e_5]=ce_1+de_2+e_4$<br>$[e_4,e_5]=fe_1+ge_2-e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,\mu,\nu)$ | $\alpha\neq 0$<br>$\mu,\nu>0$ |
| $[e_3,e_4]=ae_1+be_2+2e_5,[e_3,e_5]=ce_1+de_2-2e_4$<br>$[e_4,e_5]=fe_1+ge_2-2e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\mu,\nu,\rho)$ | $\alpha\neq 0$<br>$\mu,\nu,\rho>0$ |
| $[e_3,e_4]=ae_1+be_2+e_5,[e_3,e_5]=ce_1+de_2-e_4$<br>$[e_4,e_5]=fe_1+ge_2+e_3$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\mu,\nu,\rho)$ | $\alpha\neq 0$<br>$\mu,\nu,\rho>0$ |
| $[e_3,e_5]=ce_1+de_2-e_3$<br>$[e_4,e_5]=fe_1+ge_2-e_4$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,\mu)$ | $\alpha\neq 0$<br>$\mu>0$ |
| $[e_3,e_5]=ce_1+de_2-e_4$<br>$[e_4,e_5]=fe_1+ge_2+xe_3-2e_4$ | $\alpha e^{12}$ | There are many cases<br>See [9] | $\alpha\neq 0$ |

TAB. 7. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra

| Non vanishing Lie brackets | $r$ | Matrix of $\varrho$ |
|---|---|---|
| $[e_3,e_1]=-e_2,[e_3,e_2]=e_1,[e_4,e_1]=e_2,[e_4,e_2]=e_1$<br>$[e_5,e_1]=e_1,[e_5,e_2]=-e_2,$<br>$[e_3,e_4]=2e_5+(l_{22}-l_{21}-2l_{13})e_1-(l_{12}+l_{11}+2l_{23})e_2$<br>$[e_3,e_5]=-2e_4+(l_{23}-l_{11}+2l_{12})e_1-(l_{13}-l_{21}-2l_{22})e_2,$<br>$[e_4,e_5]=-2e_3+(l_{23}-l_{12}+2l_{11})e_1+(l_{13}+l_{22}+2l_{21})e_2$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\mu,\nu,\rho)$<br>$\mu,\nu,\rho>0$ |
| $[e_4,e_2]=ue_1,[e_5,e_1]=-\frac{a}{2}e_1,[e_5,e_2]=ve_1+\frac{a}{2}e_2,$<br>$[e_3,e_4]=xe_1+ye_2,[e_3,e_5]=be_3+ze_1+te_2,$<br>$[e_4,e_5]=ce_3+ae_4+re_1+se_2,$<br>$(a+2b)x-2tu+2yv=0,a\neq 0,b\neq 0,(3a+2b)y=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,1)$ |
| $[e_4,e_2]=ue_1,[e_5,e_1]=-\frac{a}{2}e_1,[e_5,e_2]=ve_1+\frac{a}{2}e_2,$<br>$[e_3,e_4]=xe_1,[e_3,e_5]=ze_1+te_2,$<br>$[e_4,e_5]=ae_4+re_1+se_2,a\neq 0,$<br>$ax-2tu=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\begin{pmatrix}1&\mu\\ \mu&1\end{pmatrix},1)$ |
| $[e_4,e_1]=ue_2,[e_5,e_1]=\frac{a}{2}e_1+ve_2,[e_5,e_2]=-\frac{a}{2}e_2,$<br>$[e_3,e_4]=xe_1+ye_2,[e_3,e_5]=be_3+ze_1+te_2,$<br>$[e_4,e_5]=ce_3+ae_4+re_1+se_2,$<br>$(3a+2b)x=0,a\neq 0,b\neq 0$<br>$(a+2b)y-2zu+2xv=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,1)$ |
| $[e_4,e_1]=ue_2,[e_5,e_1]=\frac{a}{2}e_1+ve_2,[e_5,e_2]=-\frac{a}{2}e_2,$<br>$[e_3,e_4]=ye_2,[e_3,e_5]=ze_1+te_2,$<br>$[e_4,e_5]=ae_4+re_1+se_2,a\neq 0,ay-2zu=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\begin{pmatrix}1&\mu\\ \mu&1\end{pmatrix},1)$ |
| $[e_4,e_1]=ue_1+upe_2,[e_4,e_2]=-\frac{u}{p}e_1-ue_2,,$<br>$[e_5,e_1]=ve_1+\frac{(2v-a)p}{2}e_2,[e_5,e_2]=-\frac{(2v+a)}{2p}e_1-ve_2$<br>$[e_3,e_4]=xe_1+ye_2,[e_3,e_5]=be_3+ze_1+te_2,a\neq 0,b\neq 0$<br>$[e_4,e_5]=ce_3+ae_4+re_1+se_2,$<br>$((2a+2b+2v)x-2zu)p-ay+2tu-2yv=0$<br>$(2xv-ax-2zu)p+(2a+2b-2v)y+2tu=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,1)$ |
| $[e_4,e_1]=ue_1+upe_2,[e_4,e_2]=-\frac{u}{p}e_1-ue_2,,$<br>$[e_5,e_1]=ve_1+\frac{(2v-a)p}{2}e_2,[e_5,e_2]=-\frac{(2v+a)}{2p}e_1-ve_2$<br>$[e_3,e_4]=xe_1+ye_2,[e_3,e_5]=ze_1+te_2,$<br>$[e_4,e_5]=ae_4+re_1+se_2,a\neq 0,b\neq 0$<br>$((2a+2v)x-2zu)p-ay+2tu-2yv=0$<br>$(2xv-ax-2zu)p+(2a-2v)y+2tu=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,\begin{pmatrix}1&\mu\\ \mu&1\end{pmatrix},1)$ |
| $[e_5,e_1]=ue_1+ve_2,[e_5,e_2]=we_1-ue_2,$<br>$[e_3,e_4]=xe_1+ye_2,[e_3,e_5]=ae_3+be_4+ze_1+te_2,$<br>$[e_4,e_5]=ce_3+de_4+re_1+se_2,$<br>$(a+d+u)x+yw=0,\ xv+(a+d-u)y=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,1)$ |
| $[e_5,e_1]=ue_1+ve_2,[e_5,e_2]=we_1-ue_2,$<br>$[e_3,e_4]=xe_1+ye_2+ae_4,[e_3,e_5]=be_4+ze_1+te_2,$<br>$[e_4,e_5]=ce_4+re_1+se_2,a\neq 0$<br>$(c+u)x-ar+yw=0$<br>$(c-u)y-as+xv=0$ | $\alpha e^{12}$ | $\mathrm{Diag}(1,1,1,1,1)$ |

TAB. 8. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra ($\alpha\neq 0$) (Continued)

| Non vanishing Lie brackets | Bivector $r$ | Matrix of $\varrho$ | Conditions |
|---|---|---|---|
| $[e_3,e_2]=xe_1-ae_4,[e_4,e_2]=ye_1+ae_3,[e_5,e_2]=ze_1$<br>$[e_3,e_5]=pe_3+qe_4+a^{-1}(-qx+py)e_1,$<br>$[e_3,e_5]=-qe_3+pe_4-a^{-1}(px+qy)e_1$ | $\alpha e^{12}$ | Diag$(1,1,1,1,1)$ | $\alpha\neq 0$<br>$a\neq 0$ |
| $[e_3,e_2]=xe_1-ae_4,[e_4,e_2]=ye_1+ae_3,[e_5,e_2]=ze_1$<br>$[e_3,e_4]=be_1$<br>$[e_3,e_5]=qe_4-a^{-1}qxe_1,$<br>$[e_3,e_5]=-qe_3-a^{-1}qye_1$ | $\alpha e^{12}$ | Diag$(1,1,1,1,1)$ | $\alpha\neq 0$<br><br>$a\neq 0, z\neq 0$ |
| $[e_3,e_2]=xe_1-ae_4,[e_4,e_2]=ye_1+ae_3,$<br>$[e_3,e_4]=be_1+ce_2$<br>$[e_3,e_5]=qe_4-a^{-1}qxe_1,$<br>$[e_3,e_5]=-qe_3-a^{-1}qye_1$ | $\alpha e^{12}$ | Diag$(1,1,1,1,1)$ | $\alpha\neq 0$<br><br>$a\neq 0$ |

TAB. 9. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra (Continued)

This theorem unknown to our knowledge can be used to build examples of Riemann-Poisson Lie algebras.

**Theorem 4.1.** *Let $(G,\langle\ ,\ \rangle)$ be an even dimensional flat Riemannian Lie group. Then there exists a left invariant differential $\Omega$ on $G$ such that $(G,\langle\ ,\ \rangle,\Omega)$ is a Kähler Lie group.*

**Proof.** Let $\mathfrak{g}$ be the Lie algebra of $G$ and $\varrho=\langle\ ,\ \rangle(e)$. According to Milnor's Theorem [11, Theorem 1.5] and its improved version [1, Theorem 3.1] the flatness of the metric on $G$ is equivalent to $[\mathfrak{g},\mathfrak{g}]$ is even dimensional abelian, $[\mathfrak{g},\mathfrak{g}]^{\perp}=\{u\in\mathfrak{g},\mathrm{ad}_u+\mathrm{ad}_u^*=0\}$ is also even dimensional abelian and $\mathfrak{g}=[\mathfrak{g},\mathfrak{g}]\oplus[\mathfrak{g},\mathfrak{g}]^{\perp}$. Moreover, the Levi-Civita product is given by

$$\mathrm{L}_a=\begin{cases}\mathrm{ad}_a & \text{if}\quad a\in[\mathfrak{g},\mathfrak{g}]^{\perp},\\ 0 & \text{if}\quad a\in[\mathfrak{g},\mathfrak{g}]\end{cases} \tag{21}$$

and there exists a basis $(e_1,f_1,\dots,e_r,f_r)$ of $[\mathfrak{g},\mathfrak{g}]$ and $\lambda_1,\dots,\lambda_r\in[\mathfrak{g},\mathfrak{g}]^{\perp}\setminus\{0\}$ such that for any $a\in[\mathfrak{g},\mathfrak{g}]^{\perp}$,

$$[a,e_i]=\lambda_i(a)f_i\quad\text{and}\quad[a,f_i]=-\lambda_i(a)e_i\,.$$

We consider a nondegenerate skew-symmetric 2-form $\omega_0$ on $[\mathfrak{g},\mathfrak{g}]^{\perp}$ and $\omega_1$ the nondegenerate skew-symmetric 2-form on $[\mathfrak{g},\mathfrak{g}]^{\perp}$ given by $\omega_1=\sum_{i=1}^{r}e_i^*\wedge f_i^*$. One can sees easily that $\omega=\omega_0\oplus\omega_1$ is a Kähler form on $\mathfrak{g}$. □

## REFERENCES


[1] Ait Haddou, M., Boucetta, M., Lebzioui, H., *Left-invariant Lorentzian flat metrics on Lie groups*, J. Lie Theory **22** (1) (2012), 269–289.

[2] Boucetta, M., *Compatibilité des structures pseudo-riemanniennes et des structures de Poisson*, C.R. Acad. Sci. Paris Sér. I **333** (2001), 763–768.

[3] Boucetta, M., *Riemann-poisson manifolds and kähler-riemann foliations*, C.R. Acad. Sci. Paris, Sér. I **336** (2003), 423–428.

[4] Boucetta, M., *Poisson manifolds with compatible pseudo-metric and pseudo-Riemannian Lie algebras*, Differential Geom. Appl. **20** (2004), 279–291.

[5] Boucetta, M., *On the Riemann-Lie algebras and Riemann-Poisson Lie groups*, J. Lie Theory **15** (1) (2005), 183–195.

[6] Deninger, C., Singhof, W., *Real polarizable hodge structures arising from foliation*, Ann. Global Anal. Geom. **21** (2002), 377–399.

[7] Dufour, J.P., Zung, N.T., *Poisson Structures and Their Normal Forms*, Progress in Mathematics, vol. 242, Birkhäuser Verlag, 2005.

[8] Fernandes, R.L., *Connections in Poisson Geometry 1: Holonomy and invariants*, J. Differential Geom. **54** (2000), 303–366.

[9] Ha, K.Y., Lee, J.B., *Left invariant metrics and curvatures on simply connected three dimensional Lie groups*, Math. Nachr. **282** (2009), 868–898.

[10] Hawkin, E., *The structure of noncommutative deformations*, J. Differential Geom. **77** (2007), 385–424.

[11] Milnor, J., *Curvatures of left invariant metrics on Lie Groups*, Adv. Math. **21** (1976), 293–329.

[12] Ovando, G., *Invariant pseudo-Kähler metrics in dimension four*, J. Lie Theory **16** (2006), 371–391.

[13] Vaisman, I., *Lectures on the Geometry of Poisson Manifolds*, Progress in Mathematics, vol. 118, Birkhäuser, Berlin, 1994.

[a] Université de Nouakchott,
Faculté des sciences et techniques
*E-mail*: pacha.ali86@gmail.com

[b] Université Cadi-Ayyad,
Faculté des sciences et techniques,
BP 549 Marrakech Maroc
*E-mail*: m.boucetta@uca.ac.ma

[c] Université de Nouakchott,
Faculté des sciences et techniques
*E-mail*: lessiadahmed@gmail.com