ARCHIVUM MATHEMATICUM (BRNO)
Tomus 61 (2025), 167–173

HYPERBOLIC SUMMATION INVOLVING THE FUNCTION $\Omega(n)$ AND LCM

Meselem Karras and Mihoub Bouderbala

Abstract. We study the sum $\sum\limits_{abc\leq x} \Omega\left([a,b,c]\right)$, where $\Omega(n)$ denotes the number of distinct prime divisors of $n \in \mathbb{Z}_{\geq 1}$ counted with multiplicity, and $[a,b,c] = \operatorname{lcm}(a,b,c)$. An asymptotic formula is derived for this sum over the hyperbolic region $\left\{(a,b,c) \in \mathbb{Z}^3_{\geq 1},\ abc \leq x\right\}$.

Introduction

Let f be an arithmetic function, and for $r \geq 2$, let

$$[n_1, n_2, \dots n_r] = \operatorname{lcm}\left(n_1, n_2, \dots n_r\right) .$$

Let $\omega\left(n\right)$ denote the number of distinct prime divisors of a positive integer $n \geq 1$, and let $\Omega\left(n\right)$ the number of prime divisors of n, counted with multiplicity. The problem of finding an asymptotic formula for sums such as

$$\sum_{n_1 n_2 \dots n_r \leq x} f\left(\gcd\left(n_1, n_2, \dots n_r\right)\right) \ \text{ or } \sum_{n_1 n_2 \dots n_r \leq x} f\left(\left[n_1, n_2, \dots n_r\right]\right)$$

where f is a suitable arithmetic function, to be speced below, has been widely studied in number theory. Previous works by researchers such as Heyman and Tóth [1] include sharp results in the two-variable case. Results for the general case are often restricted to multiplicative arithmetic functions under certain conditions, or to additive arithmetic functions for the first type of sum [2]. For the second type of sum, we first restrict our attention to the case $f = \omega^m$ with $m \geq 1$. This is, however, not our main topic in the present paper. Indeed, the case $f = \Omega^m$ where $m \geq 1$, poses substantial challenges and requires different techniques. In the present paper we are able to make a new and hopefully interesting contribution for the case $f = \Omega$, with $r = 3$.

We begin with two important results which are direct applications of two of Ivic's theorems [4].

2020 *Mathematics Subject Classification*: primary 11A05; secondary 11A25, 11N37.

Key words and phrases: prime divisors, hyperbolic summation, integer part.
Received March 21, 2025, revised November 2025. Editor C. Greither.
DOI: 10.5817/AM2025-5-167

Theorem 1. *Let $r \geq 2$ be fixed integer and N be an arbitrary fixed integer, but for which $N > r$. Then there exist computable constants $a_{r,j}$, $b_{r,j}$, $c_{r,j}$ $(a_{r,j} \neq 0)$ such that*

$$\begin{aligned}\sum_{n_1 n_2 \ldots n_r \leq x} \omega\left([n_1, n_2, \ldots n_r]\right) &= x \sum_{j=1}^{N} \left(a_{r,j} \log\log x + b_{r,j}\right) (\log x)^{r-j} \\ &\quad + x \sum_{j=r+1}^{N} c_{r,j} (\log x)^{r-j} + O\big(x (\log x)^{r-N-1}\big)\,. \end{aligned} \tag{1}$$

Theorem 2. *Let m, $N \geq 1$ and $r \geq 2$ be fixed integers. Then there exist polynomials $P_{r,m,j}\left(t\right)$ $(j = 1, 2, \ldots, N)$ of degree m in t with computable coefficients such that*

$$\begin{aligned}\sum_{n_1 n_2 \ldots n_r \leq x} \omega^m\left([n_1, n_2, \ldots n_r]\right) &= x \sum_{j=1}^{N} P_{r,m,j}\left(\log\log x\right) (\log x)^{r-j} \\ &\quad + O\big(x (\log x)^{r-N-1} \left(\log\log x\right)^m\big)\,. \end{aligned} \tag{2}$$

The authors in [1, Theorem 2.9] show the first result in the case $r = 2$ and the same result for the function $\Omega\left(n\right)$ (see Theorem 2.10).

Proof. The distinct prime divisors of the integer $[n_1, n_2, \ldots, n_r]$ are the same of the integer $n_1 n_2 \ldots n_r$. Then we obtain

$$\omega\left([n_1, n_2, \ldots, n_r]\right) = \omega\left(n_1 n_2 \ldots n_r\right)\,.$$

Therefore, for any integer $m \geq 1$

$$\begin{aligned}\sum_{n_1 n_2 \ldots n_r = n} \omega^m\left([n_1, n_2, \ldots, n_r]\right) &= \sum_{n_1 n_2 \ldots n_r = n} \omega^m\left(n\right) \\ &= \omega^m\left(n\right) \sum_{n_1 n_2 \ldots n_r = n} 1 \\ &= \omega^m\left(n\right) \tau_r\left(n\right)\,. \end{aligned}$$

Note that, for an integer $r \geq 2$,

$$\tau_r\left(n\right) = \sum_{n_1 n_2 \ldots n_r = n} 1$$

is the Piltz divisor function. Thus,

$$\begin{aligned}\sum_{n_1 n_2 \ldots n_r \leq x} \omega^m\left([n_1, n_2, \ldots, n_r]\right) &= \sum_{n \leq x} \sum_{n_1 n_2 \ldots n_r = n} \omega^m\left(n\right) \\ &= \sum_{n \leq x} \omega^m\left(n\right) \tau_r\left(n\right)\,. \end{aligned}$$

So, our results are Theorems 1 and 2 in [4]. $\square$

Theorem 3. *We have*

$$\sum_{abc\leq x} \Omega\left([a,b,c]\right) = \frac{3}{2}x\left(\log x\right)^2 \log\log x + 3(b-1)x^2\log x + \frac{(C_2-3b)}{2}x\left(\log x\right)^2 + O\left(x\log x\log\log x\right)\,,$$

where $b = A + \sum_p \frac{1}{p\,(p-1)}$ *such that* $A = \gamma + \sum_p \Big(\log\Big(1-\frac{1}{p}\Big)+\frac{1}{p}\Big)$ *and*

$$C_2 = \sum_p \frac{1}{p^3-1} \approx 0.1941\,.$$

The proof of the theorem is based on the following lemmas:

Lemma 1. *Let f be an arithmetic function. Then*

$$\sum_{abc\leq x} f\left([a,b,c]\right) = 3\sum_{an\leq x} f\left(a\right)\tau\left(n\right) - 3x\sum_{n\leq x}\frac{1}{n}\sum_{ab=n} f\Big(\gcd\left(a,b\right)\Big) + \sum_{abc\leq x} f\left(\gcd\left(a,b,c\right)\right) + O\Big(\sum_{ab\leq x} f\Big(\gcd\left(a,b\right)\Big)\Big)\,, \tag{3}$$

where $\tau\left(n\right)$ is the number of positive divisors of n.

Proof. Using the inclusion-exclusion principle, we have

$$\begin{aligned}\sum_{abc=n} f\left([a,b,c]\right) &= \sum_{abc=n} f\left(a\right) + \sum_{abc=n} f\left(b\right) + \sum_{abc=n} f\left(c\right) - \sum_{abc=n} f\Big(\gcd\left(a,b\right)\Big)\\ &\quad - \sum_{abc=n} f\Big(\gcd\left(a,c\right)\Big) - \sum_{abc=n} f\Big(\gcd\left(b,c\right)\Big) + \sum_{abc=n} f\Big(\gcd\left(a,b,c\right)\Big)\\ &= 3\sum_{abc=n} f\left(a\right) - 3\sum_{abc=n} f\Big(\gcd\left(a,b\right)\Big)\\ &\quad + \sum_{abc=n} f\Big(\gcd\left(a,b,c\right)\Big)\,.\end{aligned}$$

Thus, we obtain:

$$\begin{aligned}\sum_{abc\leq x} f\left([a,b,c]\right) &= 3\sum_{abc\leq x} f\left(a\right) - 3\sum_{abc\leq x} f\Big(\gcd\left(a,b\right)\Big) + \sum_{abc\leq x} f\Big(\gcd\left(a,b,c\right)\Big)\\ &= 3\sum_{an\leq x} f\left(a\right)\tau\left(n\right) - 3\sum_{ab\leq x} f\Big(\gcd\left(a,b\right)\Big)\sum_{c\leq \frac{x}{ab}} 1\\ &\quad + \sum_{abc\leq x} f\Big(\gcd\left(a,b,c\right)\Big)\,.\end{aligned}$$

Furthermore, since:

$$\sum_{ab\leq x} f\big(\gcd(a,b)\big) \sum_{c\leq \frac{x}{ab}} 1 = x \sum_{ab\leq x} \frac{f\big(\gcd(a,b)\big)}{ab} + O\Big(\sum_{ab\leq x} f\big(\gcd(a,b)\big)\Big)$$
$$= x \sum_{n\leq x} \frac{1}{n} \sum_{ab=n} f\big(\gcd(a,b)\big) + O\Big(\sum_{ab\leq x} f\big(\gcd(a,b)\big)\Big).$$

So, considering this last formula, we obtain the desired result. □

Lemma 2. *For $x \geq 2$, we have*

$$\sum_{an\leq x} \Omega(a)\,\tau(n) = \frac{x}{2}(\log x)^2 \log\log x + (b-1)x^2 \log x - \frac{b}{2}x(\log x)^2$$
$$+ O(x \log x \log\log x) \,, \tag{4}$$

where $b = A + \sum_p \frac{1}{p(p-1)}$ *such that* $A = \gamma + \sum_p \Big(\log\Big(1 - \frac{1}{p}\Big) + \frac{1}{p}\Big) \approx 0.2614972\ldots.$

Proof. By the well-known estimate formula,

$$\sum_{n\leq x} \tau(n) = x(\log x + C) + O\left(x^{\theta+\varepsilon}\right),$$

where $C = 2\gamma - 1$ and $\frac{1}{4} < \theta < \frac{1}{2}$, we have

$$\sum_{an\leq x} \Omega(a)\,\tau(n) = \sum_{a\leq x} \Omega(a) \sum_{n\leq \frac{x}{a}} \tau(n)$$
$$= \sum_{a\leq x} \Omega(a) \left(\frac{x}{a}\left(\log\frac{x}{a} + C\right) + O\Big(\Big(\frac{x}{a}\Big)^{\theta+\varepsilon}\Big)\right)$$
$$= x(\log x + C) \sum_{n\leq x} \frac{\Omega(n)}{n} - x \sum_{n\leq x} \frac{\Omega(n)\log n}{n} + O\left(x^{\theta+\varepsilon} \sum_{n\leq x} \frac{\Omega(n)}{n^{\theta+\varepsilon}}\right).$$

Estimate of the following sums $\sum_{n\leq x} \frac{\Omega(n)}{n}$, $\sum_{n\leq x} \frac{\Omega(n)\log n}{n}$ and $\sum_{n\leq x} \frac{\Omega(n)}{n^{\theta+\varepsilon}}$.

We know that,

$$\sum_{n\leq x} \Omega(n) = x \log\log x + bx + O\Big(\frac{x}{\log x}\Big),$$

where b is given by (4). We use a partial summation, we get

$$\sum_{n\leq x} \frac{\Omega(n)}{n} = (\log x)\log\log x + (b-1)\,x + O(\log\log x)\,.$$

Again by partial summation, we have

$$\sum_{n\leq x} \frac{\Omega(n)\log n}{n} = \log x \sum_{n\leq x} \frac{\Omega(n)}{n} - \int_2^x \frac{1}{t}\Big(\sum_{n\leq t} \frac{\Omega(n)}{n}\Big)dt$$
$$= \frac{1}{2}(\log x)^2 \log\log x + \frac{b}{2}(\log x)^2 + O\big((\log x)\log\log x\big),$$

and we have

$$\sum_{n\leq x}\frac{\Omega\left(n\right)}{n^{\theta+\varepsilon}}=O\left(x^{1-\theta-\varepsilon}\log\log x\right)\ .$$

□

Remark 4. We can use the Theorem 3 from [4] with parameters $k=3$, $m=0$ and $r=1$.

Lemma 3. *We have*

$$\sum_{n\leq x}\frac{1}{n}\sum_{ab=n}\Omega\left(\gcd\left(a,b\right)\right)=\frac{C_{\Omega}}{2}\log^{2}\left(x\right)+\left(C_{\Omega}+1\right)\log x+D_{\Omega}+O\left(x^{-1/2}\right)\ , \tag{5}$$

where C_Ω and D_Ω are two positive constants.

Proof. We apply a partial summation to the estimate (2.16) in [1], we get (5). □

Lemma 4. *We have*

$$\sum_{abc\leq x}\Omega\big(\gcd\left(a,b,c\right)\big)=\frac{C_{2}}{2}x\log^{2}\left(x\right)+O\left(x\log x\right)\ , \tag{6}$$

where $C_2=\sum_{p}\frac{1}{p^{3}-1}\approx 0.1941$.

We note that formula (6) has been proved in the general case in [2, Theorem 2.3]. Here we give the explicit form of the polynomial $P_{\Omega,2}\left(x\right)$ of degree 2. First, by Proposition 5.1 in [5], we get

$$\sum_{abc=n}\Omega\big(\gcd\left(a,b,c\right)\big)=\sum_{d^{3}m=n}\left(\mu*\Omega\right)\left(d\right)\tau_{3}\left(m\right)\ ,$$

then

$$\begin{aligned}\sum_{abc\leq x}\Omega\big(\gcd\left(a,b,c\right)\big)&=\sum_{d^{3}m\leq x}\left(\mu*\Omega\right)\left(d\right)\tau_{3}\left(m\right)\\&=\sum_{d\leq x^{1/3}}\left(\mu*\Omega\right)\left(d\right)\sum_{m\leq\left(\frac{x}{d}\right)^{1/3}}\tau_{3}\left(m\right)\ .\end{aligned}$$

We use the estimate

$$\sum_{m\leq x}\tau_{3}\left(m\right)=\frac{x\log^{2}x}{2}+O\left(x\log x\right)\ ,$$

see, e.g., Nathanson [6, Th 7.6]. According to this estimate

$$
\begin{aligned}
\sum_{abc\leq x} \Omega\big(\gcd(a,b,c)\big) &= \sum_{d\leq x^{1/3}} (\mu*\Omega)(d)\left(\frac{x}{2d^3}\log^2\left(\frac{x}{d^3}\right)+O\left(\frac{x}{d^3}\log\left(\frac{x}{d^3}\right)\right)\right)\\
&= \frac{x\log^2 x}{2}\sum_{d\leq x^{1/3}}\frac{(\mu*\Omega)(d)}{d^3}+\Big(-3x\log x+\frac{9}{2}x\Big)\\
&\quad\times\sum_{d\leq x^{1/3}}\frac{(\mu*\Omega)(d)\log d}{d^3}\\
&\quad+O\bigg(x\log x\sum_{d\leq x^{1/3}}\frac{(\mu*\Omega)(d)}{d^3}\bigg). \qquad (7)
\end{aligned}
$$

On the other hand, we have

$$
\begin{aligned}
\sum_{n=1}^{\infty}\frac{(\mu*\Omega)(n)}{n^s} &= \sum_{n=1}^{\infty}\frac{\mu(n)}{n^s}\times\sum_{n=1}^{\infty}\frac{\Omega(n)}{n^s}\\
&= \frac{1}{\zeta(s)}\times\zeta(s)\sum_p\frac{1}{p^s-1}\\
&= \sum_p\frac{1}{p^s-1},\quad \operatorname{Re}(s)>1\,.
\end{aligned}
$$

then

$$
\begin{aligned}
\sum_{d\leq x^{1/3}}\frac{(\mu*\Omega)(d)}{d^3} &= \sum_{d=1}^{\infty}\frac{(\mu*\Omega)(d)}{d^3}+O\Big(\frac{1}{x^{2/3}}\Big)\\
&= \sum_p\frac{1}{p^3-1}+O\Big(\frac{1}{x^{2/3}}\Big) = C_2+O\Big(\frac{1}{x^{2/3}}\Big)\,. \qquad (8)
\end{aligned}
$$

Using this last estimate with a partial sum, we find

$$
\sum_{d\leq x^{1/3}}\frac{(\mu*\Omega)(d)\log d}{d^3} = O(1) \qquad (9)
$$

According (8) and (9) in (7), therefore we obtain (6).

Proof of Theorem 3. By Lemma 1, we have

$$
\begin{aligned}
\sum_{abc\leq x}\Omega([a,b,c]) &= 3\sum_{an\leq x}\Omega(a)\,\tau(n)-3x\sum_{n\leq x}\frac{1}{n}\sum_{ab=n}\Omega\big(\gcd(a,b)\big)\\
&\quad+\sum_{abc\leq x}\Omega\big(\gcd(a,b,c)\big)+O\bigg(\sum_{ab\leq x}\Omega\big(\gcd(a,b)\big)\bigg),
\end{aligned}
$$

and by Lemmas 2, 3 and 4 we get

$$\sum_{abc\leq x}\Omega\left([a,b,c]\right)=\frac{3}{2}x\left(\log x\right)^{2}\log\log x+3(b-1)x^{2}\log x$$
$$+\frac{(C_{2}-3b)}{2}x\left(\log x\right)^{2}+O\big(x\log x\log\log x\big)\,,$$

which concludes the proof. □

References

[1] Heyman, R., Tóth, L., *On Certain Sums of Arithmetic Functions Involving the GCD and LCM of Two Positive Integers*, Results Math. **76** (1) (2021), 22 pp., Paper No. 49.

[2] Heyman, R., Tóth, L., *Hyperbolic summation for functions of the GCD and LCM of several integers*, Ramanujan J. **62** (1) (2022), 1–18.

[3] Iksanov, A., Marynych, A., Raschel, K., *Asymptotics of arithmetic functions of GCD and LCM of random integers in hyperbolic regions*, preprint, 2021, arXiv:2112.11892v1[math.NT].

[4] Ivic, A., *Sums of products of certain arithmetical functions*, Publ. Inst. Math. (Beograd) (N.S.) **41** (55) (1987), 31–41.

[5] Krätzel, E., Nowak, W.G., Tóth, L., *On certain arithmetic functions involving the greatest common divisor*, Cent. Eur. J. Math. **10** (2012), 761–774.

[6] Nathanson, M.B., *Elementary Methods in Number Theory*, Grad. Texts in Math., vol. 195, Springer, 2000.

University of Tissemsilt, Faculty of Science and Technology,
Algeria FIMA Laboratory,
Khemis Miliana, Algeria
E-mail: m.karras@univ-tissemsilt.dz

Khemis Miliana University (UDBKM), FIMA Laboratory,
Faculty of Matter Sciences and Computer Sciences,
Rue Thniet El Had, Khemis Miliana,
44225, Ain Defla Province, Algeria
E-mail: mihoub75bouder@gmail.com